



Impact of Adding Chitosan and Probiotic to Broiler Dietary on carcass traits*

Ahmed Othman. Hassan¹ Qanaa Hussin Ameen. Al- Jabari² Nidhal Abdulghani Mustafa³
Ahmad87.othman@gmail.com dr_qanaameen@uokirkuk.edu.iq nidhal.mustafa@su.edu.krd

^{1,2} Department of Animals Production, College of Agriculture, Kirkuk University, Kirkuk, Iraq.

³ Department of Animal Resources, College of Agriculture Engineering Sciences, University of Salahaddin, Erbil, Iraq

- Date of research received 10/07/2023 and accepted 26/07/2023.
- Part of MSc. dissertation for the first author.

Abstract

This study aims to examine the impact of adding chitosan and probiotics to the diets of broilers on the carcass characteristics of broilers. A total of 144 broiler chicks from Ross 308 were used in this study. They were of average weight (43 g). They were divided into 6 treatments with 3 replicates (8 per treatment). The first treatment was without any addition (control treatment), the second was adding 1 g/kg of probiotic, the third was adding 0.5 g/kg of chitosan, the fourth was adding 1 g/kg of chitosan, the fifth was adding 0.5 g/kg of chitosan plus 1 g/kg of probiotic and the sixth was adding 1 g/kg of chitosan plus 1 g/kg of probiotic. The foundation of this study were significant differences ($P \leq 0.05$) in the percentage of carcass refining and superiority of the third treatment, but there were no significant differences in the relative weight of the main cuttings and the relative weight of the secondary cuttings of carcass. For the internal organs (heart, gizzard and liver), there were significant differences ($p \leq 0.05$) in the relative weight of the heart, and the superiority of the fifth treatment. As for the relative weight of the gizzard, the third treatment was significantly superior ($p \leq 0.05$). The fourth treatment was significantly superior ($p \leq 0.05$) to the rest of the treatments for the relative weight of the liver. There were no significant differences in the relative weight of uneaten viscera.

Key words: chitosan, probiotic, broilers, carcass, trait.

Citation: Hasan, A., Ameen, Q., & Ghani, N. (2023). Impact of Adding Chitosan and Probiotic to Broiler Dietary on carcass traits. *Kirkuk University Journal For Agricultural Sciences*, 14(3), 75-81. doi: 10.58928/ku23.14308

Correspondence Author: Ahmed Othman Hasan- ahmad87.othman@gmail.com

Copyright: This is an open access article distributed under the terms of the creative common's attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Introduction

The yields of poultry are one of the main protein sources, they are simply digestible, also, it high in essential nutrients and minerals [1]. Poultry has been raised on a large commercial scale, and meat production is higher if we compare it with other animal products [2]. This interesting side of poultry farming agrees with the use of multiple feed additives in order to increase production, activate growth, and protect against infection with viruses and microbes [3]. Chitosan, is a non-food additive, it uses in poultry diets, it is not harmful to animal or human health; chitosan is a multiunit form of glucosamine, it makes up the lump of the external skeleton of aquatic organisms like shrimp and crabs [4,5]. Chitosan really effects on broilers' productivity performance, also, it acts as an antibacterial and antifungal agent, so it improves the health of the small intestine, improves digestion and absorption [5, 6]. Probiotics are the other and most popular food additive alternative to antibiotics [7, 8]. It basically is a microorganism this microorganism must be one of the types of poultry' intestinal bacteria [8]. It provide nutrients, aid in digestion, and constrain harmful bacteria [9]. Some reports showed that many aids could be gained by adding chitosan and probiotics to the diets of broilers. [10] The addition of chitosan to the diet of broilers was no significant differences in the relative weight of the main cuttings. Qiu et al. indicated that the addition of probiotics positively affected on the carcass characteristics [11]. In addition, Al-Salihi indicated that the addition of probiotics to the diets of broiler has a significant difference in the percentage of carcass refining but there was no significant differences in the relative weight of the main cuttings [7]. In addition, [12] found that adding chitosan to the diets of broilers showed no significant differences in the relative weight of the main cuttings.

Materials and methods

A total of 144 Ross 308 unsexed broiler chicks Kosar hatcheries provided a regular weight of 43 grams at one day of age. The chicks were randomly assigned to 6 treatment

groups, in 3 replicates (8 per group). The first treatment involved no additives (control treatment), the second treatment included 1 g/kg of probiotic, the third treatment included 0.5 g/kg of chitosan, the fourth treatment included 1 g/kg of chitosan, the fifth treatment included 0.5 g/kg of chitosan plus 1 g/kg of probiotic, and the sixth treatment included 1 g/kg of chitosan plus 1 g/kg of probiotic. Chicks were raised in separate wire cages for 35 days on a floor that was covering by sawdust 3–5 cm thick. The food (in the form of powder) was provided to the chicks and water was provided for chicks' ad libitum consumption. Two birds were taken from each treatment at the end of the trial to calculating the percentage of carcass. The live birds were considered individually by a sensitive digital scale, before the birds were slaughtered in Kosar's typical abattoir. The head, feathers and the legs were removed, and before the edible internal organs (heart, gizzard and liver) were separated. The other interior organs were also separated. The carcass was pondered individually to calculate the percentage of carcass without the eaten internal organs. The results was calculated according to the equation indicated by Becker et al. (1979) ;

The percentage of carcass% = Carcass tare weight (g) / Live body weight (g) * 100

The relative weight of the carcass parts = Carcass cut weight (g) / Carcass weight (g) * 100

The relative weight of the eaten viscera= The organ's weight (g) / Live body weight (g)*100

The relative weight of the uneaten viscera= Viscera weight (g) / Live body weight (g)*100

The relative weight of abdomen fat= Fat weight (g)/ Carcass weight (g) *100

Chitosan was imported from China, and it is a white (starch-like) substance. The German-origin Miaclost probiotic was obtained from Kosar firm.

The relative weight of visceral fat= Fat weight (g)/ Carcass weight (g) *100

These performances were calculated according to the method of [13].

mode of action of chitosan

Protects the gastric mucosa and enhances the activity of the enzyme pepsin, This is due to the nature of the membrane consisting of glycoprotein the negatively charged residue of sialic acid with the presence of positively charged chitosan, the interaction between the mucosa and chitosan leads to an increase in mucosal strength. It increases the height and density of villi and improves the internal environment of the small intestine, thus increasing the absorption of forage materials[14]

mode of action of probiotic

. It sticks to the cells lining the intestines and blocks the receptors for pathogenic bacteria, and it can make organic acids such as lactic acid, which has a toxic effect on pathogenic microorganisms which leads to improving the health status of birds and thus increasing productivity [15]

Results

The percentage of carcass %:

We note from table (2) that there was a significant superiority ($P \leq 0.05$) between T3 and T2 only, the treatment T3 was superior with T2, the highest percentage of carcass in this treatment, it was (75.11), there was no significant differences among other treatments. This study agrees with [7].

The relative weight of the carcass parts:

We note from table (3) that there were no significant differences ($P \leq 0.05$) in the relative weight of the carcass main parts (breast, drumsticks and thighs) but the highest relative weight of breast, drumsticks and thighs was the third treatment T3. We note from table (4) that there were no significant differences ($P \leq 0.05$) in the relative weight of the carcass secondary part (head neck, wings and back). This study agrees with [16].

The relative weight of the eaten viscera:

We note from table (5) that the treatment T5 significantly differences ($P \leq 0.05$) with the treatment T6 but there were no significantly

differences among other treatments regarding to the relative weight of the heart. Regarding to the relative weight of the gizzard, the treatment T3 significantly differences ($P \leq 0.05$) with the treatments T2 and T6. Also, the treatment T4 significantly differences ($P \leq 0.05$) with the other treatments that regarding to the relative weight of the liver. This study agrees with [12].

The relative weight of the uneaten viscera:

We note from table (6) that there were no significant differences ($P \leq 0.05$) in the relative weight of the abdomen fat and visceral fat, but the treatment T5 significantly differences ($P \leq 0.05$) with the other treatments regarding to the uneaten viscera. This study agrees with [17].

Table 1: Diets used in the experiment and the calculated chemical composition (kg).

Components (%)	Starterdiet (1-11 day)	Growthdiet (12-21 day)	Finisherdiet (22-42 day)
Wheat	377.25	327.25	588.85
Bran	100	100	100
Soya bean meal	320	272	165
Yellow Maize	150	200	100
Vegetarian Oil	10	10	10
Premix	10	8	--
Methionine	1.3	0.25	1
Lysine	1	1.2	2
Choline	1	1	0.5
Threonine	0.5	1.2	0.8
Enzyme	--	0.5	0.5
Anti-coccidiosis	1	0.5	0.25
Toxbond fort	--	1	1
Genex	--	--	0.5
Limestone	18.25	17	--
Calcium	8	--	--
Salt	1.6	--	--
Between finisher	--	--	7
Crude protein	22%	21%	20%
M.E.	3000	3100	3175
Energy:protein	1:15000	1:14761	1: 15875

Table 2: The effect of adding chitosan and probiotic to broiler diets on the percentage of carcass%

Treatment	live weight (g)	Carcass weight(g)	Percentage of carcass%
T1	2112.5 ±37.52 c	1552.68 ±11.30 c	73.5 ±1.50 ab
T2	2322.5 ±59.91 ab	1677.07 ±13.44 b	72.21 ±1.77 b
T3	2266.66 ±51.28 abc	1702.48 ±17.57 ab	75.11 ±1.03 a
T4	2245.83 ±37.20b c	1648.43 ±13.45 ab	73.4 ±5.36 ab
T5	2417.08 ±63.25 a	1786.22 ±18.04 a	73.9 ±1.63 ab
T6	2238.33± 28.9b c	1643.82± 13.22 ab	73.44 0.76 ab

The different letters within the same column indicates that there is a significant difference between the treatments at the level of significance $p>0.05$, values were Mean \pm standard error.

Table 3: The effect of adding chitosan and probiotic to broiler diets on the relative weight of the main carcass parts

Treatment	Breast%	Drumsticks%	Thighs%
T1	37.01 ±0.17 a	12.68 a	±0.27 13.38 ±0.32 a
T2	37.21 ±0.29 a	12.62 a	±0.12 13.08 ±0.48 a
T3	38.05 ±0.75 a	12.82 a	±0.48 13.73 ±0.51 a
T4	37.91 ±0.07 a	12.26 a	±0.01 12.65 ±0.05 a
T5	37.43 ±0.73 a	12.01 a	±0.39 13.11 ±0.74 a
T6	37.33 ± 0.79 a	12.64 ±0.11 a	12.96 ±0.18 a

The different letters within the same column indicates that there is a significant difference between the treatments at the level of significance $p>0.05$, values were Mean \pm standard error.

Table 4: The effect of adding chitosan and probiotic to broiler diets on the relative weight of the secondary carcass parts

Treatment	Head-neck%	Wings%	Back%
T1	3.89 ±0.11 a	9.81 ±0.27 a	21.25 ±0.59 a
T2	4.6 ±0.24 a	9.69 ±0.27 a	21.35 ±1.27 a
T3	4.33 ±0.24 a	10.1 ±0.39 a	21.94 ±0.95 a
T4	4.58 ±0.09 a	9.99 ±0.19 a	21.65 ±0.08 a
T5	4.83 ±0.37 a	10.11 ±0.14 a	21.35 ±0.13 a
T6	4.41 ±0.16 a	10.3 ±0.15 a	21.91 ±0.59 a

The different letters within the same column indicates that there is a significant difference between the treatments at the level of significance $p>0.05$, values were Mean \pm standard error.

Table 5: The effect of adding chitosan and probiotic to broiler diets on the relative weight of the eaten viscera

Treatment	Heart	Gizzard	Liver
T1	0.02 ±0.02 ab	1.14 ±0.05 b	2.28 ±0.07 b
T2	0.55 ±0.04 ab	1.17 ±0.07 b	2.12 ±0.02 c
T3	0.61 ±0.03 ab	1.32 ±0.12 a	2.28 ±0.14 b
T4	0.55 ±0.02 ab	1.2 ±0.07 ab	2.41 ±0.12 a
T5	0.62 ±0.03 a	1.2 ±0.02 ab	2.28 ±0.06 b
T6	0.51 ±0.02 b	1.03 ±0.02 c	2.28 ±0.13 b

The different letters within the same column indicates that there is a significant difference between the treatments at the level of significance $p>0.05$, values were Mean \pm standard error.

Table 6: The effect of adding chitosan and probiotic to broiler diets on the relative weight of the uneaten viscera

Treatment	Abdominal fat	Viscera fat	Uneaten viscera
T1	1.04 ±0.16 a	0.18 ±0.02a	6.24 ±0.19a
T2	0.88 ±0.20 a	0.11 ±0.03a	6.17 ±0.46a
T3	0.64 ±0.07 a	0.13 ±0.01a	6.21 ±0.65a
T4	0.53 ±0.06 a	0.1 ±0.01a	6.21 ±0.28 a
T5	0.78 ±0.02 a	0.16 ±0.01a	6.27 ±0.28a
T6	0.77 ±0.03 a	0.15 ±0.03 a	6.19 ±0.56 a

The different letters within the same column indicates that there is a significant difference between the treatments at the level of significance $p>0.05$, values were Mean \pm standard error.

discussion

The significant superiority that appeared in table (2) may be due to the increased utilization of protein by increasing the secretion of the enzyme trypsin in the stomach and thus increasing the utilization of amino acids [16]. The reason for the results in the table (3) and the table (4) may be due to that chitosan works to reduce the digestion of fats in the diet. chitosan increases the viscosity of dry organic matter that must be digested, as well as binds fats, preventing them from digestion and absorption in the intestine. and this effect leads to a reduction in fat deposition in the parts [12]. According to the table (5) The reason for obtaining this result may be due to the increase

in the size of the liver as a result of the increase in immunity due to the use of chitosan, which increased the size of the liver within the normal limits, and this increase in weight may be due to the increase in the fat content in the liver, as for the weight of the heart and gizzard it may be due to chitosan, as it is known to increase weight in general, which positively affects the weight of the internal organs [12]. The results of the table (6) due to the fact that chitosan works to reduce the digestion and absorption of fats by binding to fat molecules and converting them into high-viscosity compounds, which leads to impeding the digestion of these fats, and therefore they are excreted outside the body, which leads to a decrease in body fat [18].

Conclusion

The addition of chitosan and probiotics to the diets of broilers caused a significantly ($P < 0.05$) favorable effect. This effect is demonstrated by increased carcass traits. In my opinion T3 and T5 achieved the best result.

Acknowledgments

This study was conducted at animal production farms at Kosar Agricultural Research Company in Arbil. Great thanks to the staff in these Feld for providing the equipment, requirements, and facilities.

References

- [1] Windhorst H W. Changes in poultry production and trade worldwide. *World's Poultry Sci. J.* 2013: 62: 585-602.
- [2] Al-Bazy F I, Abdulateef S M, and Sulimn B F. Impact of feeds containing optifeed® vëo® premium, and oleobiotec® on the lipid peroxidation of male broilers under heat stress. *J. of Life Science and Applied Research.* 2022: 3:25-31.
- [3] Kareem B S, and Al-Dalawi R H.. Effect of adding L., Carnitine with herbal methionine and sunflower seed oil on the biochemical characteristics of broiler's blood. *J. of Kirkuk University for Agricultural Sciences.* 2022: 13(4).
- [4] Jasim H H, and Nafea H H.. Effect of chitosan and antibiotic adding to corn-soybean diet on the productive performance of broiler chickens. *Indian J. of Ecology.* 2021:48:10-14.
- [5] Friedman M, and Juneja V K. Review of antimicrobial and antioxidative activities of chitosans in food. *J. of Food Protection.* 2010: 73:1737-1761.
- [6] Singla A K, and Chawla M. 2001. Chitosan: Some pharmaceutical and biological aspects-an update. *J. of Pharmacy and Pharmacology.* 2001: 53:1047-1067.
- [7] Al-Salihy S A, and Hussaini M I A N A. Effect of different levels of boswellia plant extract in drinking water (Photovoltaic Catalyst) and the bio probiotic in the diet on the growth characteristics, physical characteristics and blood biochemistry of quail bird. *J. of Kirkuk University for Agricultural Sciences,* 2020: 11(3).
- [8] Lutful Kabir S M. The role of probiotics in the poultry industry. *International J. of Molecular Sciences.* 2009: 10:3531-3546.
- [9] Al-Sakr S N A, and AL-Neemi M I A. The effect of adding levels of the, amino acid threonine (Thr) in the low protein diets upon the carcass traits, Some physiological and bio chemical parameters of the broiler blood. *J. of Kirkuk University for Agricultural Sciences.* 2022:13(4).
- [10] Nuengjamnong C, and Angkanaporn K. Efficacy of dietary chitosan on growth performance, haematological parameters and gut function in broilers. *Italian J. of Animal Science.* 2018: 17:428-435.
- [11] Qiu K, Wang X, Zhang H, Wang J, Qi G, and Wu S. Dietary supplementation of a new probiotic compound improves the growth performance and health of broilers by altering the composition of cecal microflora. *Biology.* 2022: 11:633.
- [12] Tufan, T., and C. Arslan . Dietary supplementation with chitosan oligosaccharide affects serum lipids and nutrient digestibility in broilers. *South African J. of Animal Science.* 2020: 50:5.
- [13] BECKER, W. A., SPENCER, J. V., MIROSH, L. W., & VERSTRATE, J. A. Prediction of fat and fat free live weight in broiler chickens using backskin fat, abdominal fat, and live body weight. *Poultry Science.* 1979: 58(4), 835-842.
- [14] Ding, X.L.. Effects of Chitosan On Growth performance In Broiler Chickens And The Underlying Growth-Stimulating Mechanism. Abstract . Thesis.Master 2006 Animal Nutrition and Feed Sci. China. <http://www.globethesis>
- [15] Abdulwahid S. A. Effect of adding probiotic (BioSB- Gold) to the dietary on some productive performance and carcass cut of proiler chickens. *Al-Forat J. for Agricultural science.* 2017: 744-751.
- [16] Khoobani, M., Hasheminezhad, S. H., Javandel, F., Nosrati, M., Seidavi, A., Kadim, I. T., Kadim and Tufarelli, V. Effects of dietary chicory (*Chicorium intybus* L.) and probiotic blend as natural feed additives on performance traits, blood biochemistry, and gut microbiota of broiler chickens. *Antibiotics.* 2019: 9(1): 5.
- [17] Abdel-Baset Sh., Ashour E. A., El-Hack M. E. and El-Mekkawy M. M. Effect of different levels of pomegranate peel powder and probiotic supplementation on growth, carcass traits, blood

serum metabolites, antioxidant status and meat quality of broilers. *Ani.Biotec.* 2020: 33(4): 690-700.
[18] Lokman, I. H., E. B. Ibitoye, M. N. M. Hezmee, Y. M. Goh, A. B. Z. Zuki, and A. A. Jimoh. Effects of chitin and chitosan from cricket and shrimp on

growth and carcass performance of broiler chickens. *Tropical animal health and production.* 2019:51(8)2219-2225.



تأثير اضافة الكايتوسان والمعزز الحيوي الى علائق فروج اللحم على صفات الذبيحة

احمد عثمان حسن قانع حسين امين الجباري نضال عبدالغني مصطفى
Ahmad87.othman@gmail.com dr_qanaameen@uokirkuk.edu.iq nidhal.mustafa@su.edu.krd

- ¹ قسم الانتاج الحيواني، كلية الزراعة، جامعة كركوك، كركوك، العراق.
³ قسم الثروة الحيوانية، كلية علوم الهندسة الزراعية، جامعة صلاح الدين- اربيل، العراق.
• تاريخ استلام البحث 10/07/2023 وتاريخ قبوله 26/07/2023
• البحث مستل من رسالة ماجستير للباحث الاول .

الملخص

الهدف من هذه الدراسة معرفة تأثير اضافة الكايتوسان chitosan والمعزز الحيوي الى عليقة فروج اللحم على صفات الذبيحة. وفي هذه التجربة تم استخدام 144 فرخة من سلالة Ross 308 غير مجنسة والتي تم تجهيزها من مفاقس شركة كوسار وكانت بوزن ابتدائي (43غم) موزعة على ستة معاملات وكل معاملة بثلاث مكررات بواقع ثمانية افراخ لكل مكرر. وكانت المعاملات كمايلي: المعاملة الاولى: عليقة السيطرة القياسية، والمعاملة الثانية عليقة قياسية مع اضافة 1غم من المعزز الحيوي/كغم من العلف، والمعاملة الثالثة: عليقة قياسية مع اضافة 0,5 غم من الكايتوسان/ كغم من العلف، المعاملة الرابعة: عليقة قياسية مع اضافة 1غم من الكايتوسان/ كغم من العلف، المعاملة الخامسة: عليقة قياسية مع اضافة 0,5غم من الكايتوسان/كغم من العلف بالاضافى الى 1غم من المعززالحيوي/ كغم من العلف، المعاملة السادسة: عليقة قياسية مع اضافة 1غم من الكايتوسان/كغم من العلف بالاضافة الى 1غم من المعزز الحيوي/كغم من العلف. ظهرت فروقات معنوية ($p \leq 0.05$) في نسبة التصافي للذبيحة وبتفوق المعاملة الثالثة بينما لم تظهر فروقات معنوية في الوزن النسبي للقطيعات الرئيسية (الصدر، عصا الطبال، وقطعة الفخذ) وكذلك لم تظهر فروقات معنوية في الوزن النسبي للقطيعات الثانوية (الرقبة ، الاجنحة والظهر). اما بالنسبة للاحشاء الداخلية المأكولة (القلب ، القانصة والكبد) فقد ظهرت فروقات معنوية ($p \leq 0.05$) في الوزن النسبي للقلب بتفوق المعاملة الخامسة اما في الوزن النسبي للقانصة فقد تفوقت المعاملة الثالثة اما الوزن النسبي للكبد فقد تفوقت المعاملة الرابعة. اما في صفة الوزن النسبي للاحشاء الداخلية غير المأكولة (دهن البطن، دهن الاحشاء والاحشاء غير المأكولة) فلم تظهر فروقات معنوية في الوزن النسبي للاحشاء غير المأكولة.

الكلمات المفتاحية: الكايتوسان، المعزز الحيوي، فروج اللحم، الذبيحة، صفات.