



Comparison the Effect of Adding Local *Anemone coronaria* L. Flower Powder and Vitamin C in the diet on the Productive Performance of Laying Hens Exposed to Heat Stress in Summer

Mohammed Fadhil Abed Al-Khayyat¹
mfkyatt@gmail.com

Ammar Qahtan Shanoon²
ammar.qahtan@uokirkuk.edu.iq

¹ Ministry of Environment, Baghdad, IRAQ.

² Department of Animal production, College of Agriculture, University of Kirkuk, Kirkuk, IRAQ.

- Part of M.Sc. Dissertation for the first author.
- Date of research received 3/2/2024 and accepted 2/3/2024.

Abstract

The effect of adding powdered local *Anemone coronaria* L. flowers and vitamin C on the productive performance of laying hens raised in the summer and exposed to heat stress was studied. The experiment was completed using (120) Lohmann Brown birds at (30) weeks old, during July, August, and September, for a period of (84) days, which included a 14-day preparatory period. The experiment period was divided into (5) productive periods, where the birds were randomly distributed into (6) treatments with (5) replicates, and for each replicate (4) birds, where the treatment was T1: a control Diet-free of any additives, T2: a control diet with vitamin C added. At a rate of 250 mg/kg of feed, T3: a control diet to which *Anemone coronaria* L. flower powder is added. At a rate of 2 g/kg of feed, T4: a control diet to which *Anemone coronaria* L. flower powder is added, at a rate of 4 g/kg of feed, T5: a control diet with the addition of *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed and Vitamin C at a rate of 250 mg/kg feed, T6: a control diet with the addition of *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed and Vitamin C at a rate of 250 mg/ kg of feed. The results of the field experiment showed that there were significant differences in the general average of productive periods at the level ($P \leq 0.05$) in all the studied traits. Treatment T6 recorded the highest percentage of egg production, and the highest average egg weight in treatments T5 and T3 respectively, and the highest average egg mass was in treatment T6, and the highest rate of feed consumption was in treatment T2, the best food conversion factor was in the treatments T5 and T6 respectively. The control treatment T1 witnessed the highest mortality rate with a decrease in the average live weight of birds. It is noted from the results of the experiment that the additions of *Anemone coronaria* L. flower powder and vitamin C led to A reducing the effects of heat stress that affects birds when exposed to high temperatures in summer, and thus the positive effect was clear in improving the productive performance and health of the birds.

Keywords: *Anemone coronaria* L., Vitamin C, productive performance, Laying hens, Heat stress.

Citation: Al-Khayyat, M., & Shanoon, A. (2024). Comparison the Effect of Adding Local *Anemone coronaria* L. Flower Powder and Vitamin C in the diet on the Productive Performance of Laying Hens Exposed to Heat Stress in Summer. *Kirkuk University Journal for Agricultural Sciences*, 15(1), 151-163. doi: 10.58928/ku24.15115

Correspondence Author: Mohammed Fadhil Abed Al-Khayyat - mfkyatt@gmail.com

Copyright: This is an open access article distributed under the terms of the creative common's

License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Introduction

The poultry industry contributes significantly as a suitable source for achieving global food security, as it can transform a wide range of agricultural by-products into high-energy, protein-rich meat and eggs and other essential nutrients for humans while improving economic conditions [1]. Climate change and high temperatures pose a serious threat to animal production and are among the seasonal problems faced by breeders in Iraq, where poultry are exposed to heat stress, which negatively affects productive performance, health, and high mortality rates [2,3]. Heat stress leads to weight loss, decreased egg production, and even the death of birds, and its impact on poultry breeding and production is global and brings huge financial losses [4]. High temperatures in Iraq in the summer are one of the most seasonal problems that affect poultry production, as many breeders stop production in addition to the losses they suffer [2].

Nutritional supplements are the most feasible approach to overcome the effects of heat stress and improve productive performance, as many additives are used, such as vitamins, minerals, and herbs [5]. The *Anemone coronaria* L. plant is considered one of the medicinal plants of great importance because its various parts contain one or several active substances with biological effects in treating diseases, whether in their natural form or when extracted [6,7]. The extract of *Anemone coronaria* L. flowers also improves production performance, is antioxidant, analgesic, and antipyretic, and enhances the health of birds exposed to stress, thus reducing the rate of economic losses resulting from weight loss or deaths due to heat stress [8,9].

Therefore, this study, which is considered the first of its kind in Iraq and the world, aims to use the powdered flowers of the local *Anemone coronaria* L. plant as a safe feed additive and to know its effect on the productive traits of poultry birds exposed to heat stress while reducing the economic losses facing breeders in the summer and improving production performance by relying on Available natural sources, comparing them to

vitamin C, and studying the changes occurring in production characteristics.

Materials and methods

The experiment was conducted in the summer for (84) days, divided into five productive periods (14) days/period in order to measure the characteristics studied for the experiment. It was preceded by a (14) day preparatory period. The experiment was completed using (120) Lohmann Brown birds at the age of (30) weeks. They were placed in aluminum cages consisting of four floors vertically. They were placed in cages consisting of four floors vertically. The cages contained automatic plastic feeders and nipples. Table (1) shows the components of the feed and the chemical analysis. Based on NRC 1994 [10]. A lighting system of (16) hours of light and (8) hours of darkness was used, according to the recommendations of the breeding manual. The treatments were as follows: Treatment T1: a control diet free of any additives, T2: a control diet with vitamin C added. At a rate of 250 mg/kg of feed, T3: a control diet to which *Anemone coronaria* L. flower powder is added. At a rate of 2 g/kg of feed, T4: a control diet to which *Anemone coronaria* L. flower powder is added, at a rate of 4 g/kg of feed, T5: a control diet with the addition of *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed and Vitamin C at a rate of 250 mg/kg feed, T6: a control diet with the addition of *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed and Vitamin C at a rate of 250 mg/ kg of feed.

| Feed ingredients | Percentage of feed ingredients |
|------------------|--------------------------------|
| Wheat | 16.32 |
| Corn | 40.5 |
| Oil | 1 |
| Barley | 4 |
| Soybean meal 48% | 25.7 |
| Laymix-2.5 | 2.5 |
| Lysine | 0.01 |
| Methionine | 0.07 |
| Limestone | 9 |
| T. Salt | 0.2 |
| Choline chloride | 0.25 |
| DCP | 0.45 |
| Total | %100 |

| Amount | chemical composition |
|-------------------------|----------------------|
| Energy Kcal/kg | 2708 |
| Protein % | 18.37 |
| Lysine % | 1 |
| Methionine | 0.47 |
| Methionine & Cysteine % | 0.69 |
| Ca % | 4.38 |
| p % | 0.6 |
| ME/CP Ratio | 167.03 |

Statistical analysis

A completely randomized design (CRD) was used to study the effect of parameters on the studied traits using the statistical program (SAS), and the significant differences between the means were tested using the Duncan multilevel test [11], according to the following mathematical model: $Y_{ij} = \mu + T_i + e_{ij}$

Results

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the average live body weight: The results are shown in Table (2). There are significant

differences ($P \leq 0.05$) in the average live body weight of the birds at the end of the experiment compared to their weights at the beginning of the experiment, where the treatment was recorded. T4 had the highest weight gain rate, which amounted to (96.3) g, followed by treatment T6 (83.68) g, treatment T3 (38) g, treatment T5 (25.36) g, then treatment T2 recorded (21.81) g, with increase rates of 5.82%, 5.05%, 2.29%, 1.53% and 1.32%, respectively, compared to treatment T1, which witnessed an average weight decrease of (-51.35) g, meaning -3.1% from the weight of the birds at the beginning of the experiment.

Table (2) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the live body weight gain (g) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treatment | Weight gain rate/g | Percentage of weight gain % |
|-----------|-------------------------|-----------------------------|
| T1 | -51.35 \pm 25.71 c | -3.10 \pm 1.55 c |
| T2 | 21.81 \pm 20.20 b | 1.32 \pm 1.22 b |
| T3 | 38.00 \pm 26.76 ab | 2.29 \pm 1.62 ab |
| T4 | 96.30 \pm 23.61 a | 5.82 \pm 1.43 a |
| T5 | 25.36 \pm 21.26 ab | 1.53 \pm 1.28 ab |
| T6 | 83.68 \pm 17.53 ab | 5.05 \pm 1.06 ab |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the percentage of egg production (H.D.%): The results in Table (3) show that there are no significant differences in the percentage of egg production between the treatments in the first, second, third, and fourth production periods, and there are significant differences ($P \leq 0.05$) in the fifth productive period, where all treatments, without significant differences between them, outperformed the control treatment, as

treatment T3 recorded (94.29%), treatment T6 (93.57%), treatments T5 and T2 (92.5%), and treatment T4 (89.94%) compared to treatment T1 (82.38%), which is the lowest recorded level. The general rate of production periods also witnessed significant differences, as treatment T6 recorded the highest production rate of (92.29%), followed by treatment T4 (89.64%), treatment T5 (88.64%), treatment T3 (88.61%), then treatment T1 (87.88%) and treatment T2. (86.34%).

Table (3) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the percentage of egg production (H.D.%) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Egg production at different productive period | | | | | Average of egg production |
|--------|---|-----------------------|-----------------------|-----------------------|-----------------------|---------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| T1 | 85.44 \pm 2.73 a | 93.93 \pm 3.33 a | 90.12 \pm 2.89 a | 87.54 \pm 4.32 a | 82.38 \pm 2.88 b | 87.88 \pm 1.56 b |
| T2 | 80.71 \pm 3.45 a | 86.07 \pm 1.18 a | 86.07 \pm 2.85 a | 86.34 \pm 4.01 a | 92.50 \pm 1.91 a | 86.34 \pm 1.40 b |
| T3 | 81.58 \pm 2.82 a | 86.37 \pm 4.03 a | 87.86 \pm 3.31 a | 92.98 \pm 2.35 a | 94.29 \pm 2.07 a | 88.61 \pm 1.54 ab |
| T4 | 86.07 \pm 2.49 a | 91.43 \pm 2.42 a | 90.00 \pm 2.80 a | 91.07 \pm 3.29 a | 89.64 \pm 3.93 a | 89.64 \pm 1.30 ab |
| T5 | 83.57 \pm 2.49 a | 89.64 \pm 1.43 a | 89.64 \pm 2.07 a | 87.86 \pm 4.28 a | 92.50 \pm 1.73 a | 88.64 \pm 1.22 ab |
| T6 | 86.79 \pm 1.45 a | 91.07 \pm 1.26 a | 94.64 \pm 1.26 a | 95.36 \pm 2.50 a | 93.57 \pm 1.34 a | 92.29 \pm 0.92 a |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the average egg weight: The results in Table (4) show that there are no significant differences in the average egg weight between the treatments in the first, second, fourth, and fifth production periods, and there are significant differences ($P \leq 0.05$). In the third productive period, treatment T5 recorded the highest average egg weight of (60.2) g,

followed by treatment T3 (59.5) g, treatment T6 (59.19) g, treatment T4 (58.82) g, treatment T1 (58.23) g, and then treatment T2 (57.45) g. The general rate of production periods also witnessed significant differences, as the highest weight rate was recorded in treatment T5 (60.09) g and treatment T3 (59.84) g, followed by treatment T6 (59.63) g and treatment T4 (59.13) g, then treatment T1 (58.44) g and treatment T2. (58.25) g.

Table (4) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the average egg weight (g) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Egg weight at different Productive periods | | | | | Average of egg weight |
|--------|--|-----------------------|------------------------|-----------------------|-----------------------|------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| T1 | 57.66 \pm 0.86 a | 57.72 \pm 1.19 a | 58.23 \pm 1.21 ab | 59.28 \pm 0.94 a | 59.33 \pm 0.68 a | 58.44 \pm 0.43 b |
| T2 | 57.07 \pm 0.64 a | 56.13 \pm 0.83 a | 57.45 \pm 0.80 b | 59.67 \pm 1.12 a | 60.94 \pm 1.35 a | 58.25 \pm 0.54 b |
| T3 | 59.04 \pm 1.36 a | 58.30 \pm 0.66 a | 59.50 \pm 0.80 ab | 61.06 \pm 1.10 a | 61.30 \pm 0.68 a | 59.84 \pm 0.46 a |
| T4 | 58.35 \pm 0.80 a | 58.76 \pm 1.02 a | 58.82 \pm 0.79 ab | 59.72 \pm 0.95 a | 60.03 \pm 0.93 a | 59.13 \pm 0.39 ab |
| T5 | 59.50 \pm 0.81 a | 58.43 \pm 0.48 a | 60.20 \pm 0.70 a | 60.38 \pm 1.07 a | 61.95 \pm 0.46 a | 60.09 \pm 0.38 a |
| T6 | 56.70 \pm 0.58 a | 57.64 \pm 0.69 a | 59.19 \pm 0.50 ab | 61.27 \pm 0.71 a | 62.00 \pm 0.73 a | 59.36 \pm 0.49 ab |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the average egg mass: The results in Table (5) show that there are no significant differences in the average egg weight in the first and fourth productive periods, and there are significant differences ($P \leq 0.05$) in the second productive period. The third and fifth: in the second production period, treatment T1 recorded the highest rate of (54.26) g, and in the third production period, treatment T6

recorded the highest rate of (56.01) g. In the fifth production period, all treatments outperformed the control treatment T1, which recorded the lowest level of (48.8) g. The general rate of production periods also witnessed significant differences, as the highest rate was recorded in treatment T6 (54.83) g, followed by treatment T5 (53.28) g, treatment T3 (53.07) g, treatment T4 (53.04) g, then treatment T1 (51.34) g and treatment T2 (50.34) g.

Table (5) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the average egg mass (g/ bird/ day) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Egg mass at different Productive periods | | | | | Average of egg mass |
|--------|--|------------------------|------------------------|-----------------------|-----------------------|------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| T1 | 49.26 \pm 1.77 a | 54.26 \pm 2.50 a | 52.48 \pm 2.02 ab | 51.90 \pm 2.72 a | 48.80 \pm 1.21 b | 51.34 \pm 0.96 b |
| T2 | 46.02 \pm 1.74 a | 48.30 \pm 0.73 b | 49.49 \pm 2.06 b | 51.57 \pm 2.82 a | 56.33 \pm 1.34 a | 50.34 \pm 1.05 b |
| T3 | 48.08 \pm 1.37 a | 50.31 \pm 2.19 ab | 52.30 \pm 2.28 ab | 56.81 \pm 2.06 a | 57.82 \pm 1.62 a | 53.07 \pm 1.10 ab |
| T4 | 50.27 \pm 1.87 a | 53.79 \pm 2.19 ab | 53.00 \pm 2.20 ab | 54.36 \pm 2.01 a | 53.78 \pm 2.39 a | 53.04 \pm 0.92 ab |
| T5 | 49.75 \pm 1.82 a | 52.38 \pm 1.03 ab | 53.92 \pm 0.98 ab | 53.06 \pm 2.75 a | 57.29 \pm 0.84 a | 53.28 \pm 0.84 ab |
| T6 | 49.19 \pm 0.74 a | 52.48 \pm 0.71 ab | 56.01 \pm 0.80 a | 58.41 \pm 1.60 a | 58.05 \pm 1.46 a | 54.83 \pm 0.85 a |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the rate of feed consumption: The results are shown in Table (6). There are significant differences ($P \leq 0.05$) in all production periods. In the first production period, treatment T2 (98.58) g was significantly superior, and in the second period, treatment T2 (99.05) g was superior to treatment T1 (98.41) g, and in the third production period, treatment T2 (99.69) g was significantly superior, and in the fourth and fifth production periods, all The

coefficients were significantly compared to the control treatment T1, which recorded the lowest consumption rate of (93.58) g and (95.09) g/ feed/day, respectively. The general rate of production periods also witnessed significant differences, as the highest rate was recorded for treatment T2 (99.08) g, followed by treatment T6 (97.33) g, then treatment T4 (96.85) g, treatment T1 (96.43) g, then treatment T3 (95.05) g, and then treatment T5 (93.22) g.

Table (6) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the rate of feed consumption (g feed/ day) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Feed consumption at different Productive periods | | | | | Average of feed consumption |
|--------|--|------------------------|------------------------|-----------------------|-----------------------|-----------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| T1 | 96.13 \pm 1.19 ab | 98.41 \pm 0.51 a | 98.65 \pm 0.70 ab | 93.85 \pm 1.10 b | 95.09 \pm 1.08 b | 96.43 \pm 0.54 bc |
| T2 | 98.58 \pm 0.69 a | 99.05 \pm 0.59 a | 99.69 \pm 0.31 a | 98.53 \pm 0.60 a | 99.53 \pm 0.47 a | 99.08 \pm 0.24 a |
| T3 | 90.04 \pm 1.42 c | 92.87 \pm 1.43 b | 96.28 \pm 1.12 b | 98.57 \pm 0.74 a | 97.47 \pm 1.26 a | 95.05 \pm 0.81 c |
| T4 | 94.78 \pm 0.86 b | 94.08 \pm 0.69 b | 97.66 \pm 0.13 ab | 98.66 \pm 0.41 a | 99.08 \pm 0.39 a | 96.85 \pm 0.47 bc |
| T5 | 90.82 \pm 0.48 c | 86.76 \pm 2.05 c | 92.70 \pm 1.96 c | 96.79 \pm 0.82 a | 99.03 \pm 0.37 a | 93.22 \pm 1.04 d |
| T6 | 95.64 \pm 0.39 b | 95.69 \pm 0.66 ab | 98.17 \pm 0.41 ab | 97.71 \pm 0.40 a | 99.45 \pm 0.18 a | 97.33 \pm 0.35 ab |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the feed conversion factor: The results in Table (7) show that there are no significant differences in the fourth production period, but there are significant differences ($P \leq 0.05$) in the rest of the periods. In the first, second, and third production periods, treatment T2 was significantly superior to the rest of the transactions, recording (1.96, 1.83, and 1.89), respectively, and in the fifth production period,

treatment T1 was significantly superior to the rest of the transactions, recording (1.95). The general rate of production periods also witnessed significant differences, as the highest food conversion factor was recorded in treatment T2 (1.99), then treatment T1 (1.89), followed by treatment T4 (1.84) and treatment T3 (1.81), and the lowest food conversion factor was for treatment T6 (1.78) and treatment T5. (1.76).

Table (7) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the feed conversion factor (g feed/g eggs) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Feed conversion factor at different Productive periods | | | | | Average of FCR |
|--------|--|----------------------|-----------------------|----------------------|-----------------------|-----------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| T1 | 1.96 \pm 0.07 ab | 1.83 \pm 0.08 b | 1.89 \pm 0.07 ab | 1.83 \pm 0.12 a | 1.95 \pm 0.06 a | 1.89 \pm 0.04 \pm ab |
| T2 | 2.16 \pm 0.09 a | 2.05 \pm 0.02 a | 2.03 \pm 0.08 a | 1.94 \pm 0.12 a | 1.77 \pm 0.05 b | 1.99 \pm 0.04 \pm a |
| T3 | 1.88 \pm 0.06 b | 1.86 \pm 0.08 b | 1.86 \pm 0.09 ab | 1.75 \pm 0.08 a | 1.69 \pm 0.07 b | 1.81 \pm 0.03 \pm bc |
| T4 | 1.90 \pm 0.08 b | 1.76 \pm 0.08 b | 1.86 \pm 0.08 ab | 1.83 \pm 0.07 a | 1.86 \pm 0.08 ab | 1.84 \pm 0.03 \pm bc |
| T5 | 1.84 \pm 0.07 b | 1.66 \pm 0.05 b | 1.72 \pm 0.06 b | 1.85 \pm 0.10 a | 1.73 \pm 0.03 b | 1.76 \pm 0.03 \pm c |
| T6 | 1.95 \pm 0.03 ab | 1.83 \pm 0.03 b | 1.75 \pm 0.03 b | 1.68 \pm 0.04 a | 1.72 \pm 0.04 b | 1.78 \pm 0.02 \pm c |

* Different letters within column indicate the presence of significant differences ($P \leq 0.05$) between the treatments.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg Feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

The effect of adding the local *Anemone coronaria* L. flower powder and vitamin C on the mortality rate: The results in Table (8) show the number of bird deaths in each treatment and the productive period during which the death occurred. We note that the highest number of birds dying was for treatment T1, where 4 birds out of 20 birds were distributed as follows: 3 birds in the first productive period and one bird in the fifth and final productive period, meaning a percentage of 20% of the birds lost in treatment T1,

followed by treatment T3. The number of perishing birds was 2 out of 20 birds, distributed as follows: one bird in the first productive period and one bird in the second productive period, meaning a mortality rate of 10% of the total birds in treatment T3. As for the rest of the treatments, their birds did not witness any death throughout the experimental period of 70 days.

Table (8) Comparison of the effect of adding local *Anemone coronaria* L. flower powder and vitamin C in the diet on the mortality (%) of laying hens exposed to heat stress in summer (mean \pm standard error)

| Treat. | Mortality % at different Productive periods | | | | | Total number of mortality | Percentage of mortality |
|--------|---|---|---|---|---|---------------------------|-------------------------|
| | 1 | 2 | 3 | 4 | 5 | | |
| T1 | 3 | 0 | 0 | 0 | 1 | 4 | 20 % |
| T2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| T3 | 1 | 1 | 0 | 0 | 0 | 2 | 10 % |
| T4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| T5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| T6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |

* The numbers inside the columns represent the number of birds that died during production periods.

** The duration of one production period is two weeks.

*** Treatments: T1: control without any addition, T2: add vitamin C at a rate of 250 mg/kg feed, T3: add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed, T4: add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed, T5: Add *Anemone coronaria* L. flower powder at a rate of 2 g/kg feed + Vitamin C at a rate of 250 mg/kg feed, T6: Add *Anemone coronaria* L. flower powder at a rate of 4 g/kg feed + Vitamin C at a rate of 250 mg/kg feed.

Discussion

The positive effect of adding *Anemone coronaria* L. flower powder and vitamin C was clear on the productive traits studied in the experiment, represented by (egg production rate, egg weight rate, egg mass rate, feed consumption rate, feed conversion factor, and mortality rate). The results regarding the average live weight of the birds at the end of the experiment compared to their weights at the beginning of the experiment show that the only treatment that caused the birds to lose their average weight was the T1 treatment. This may be due to the powder of *Anemone coronaria* L. flowers containing active substances such as saponins and essential oils, which can be considered new factors. Important in enhancing growth performance, the effect could be related to its improved effect on intestinal growth and increased protease activity, which leads to increased protein digestion and absorption of nutrients [12]. It also promotes growth and improves immunity [13].

The studied productive characteristics related to eggs, such as the egg production percentage, which achieved its highest significant level in treatment T6, amounting to (92.29%) and the average egg weight, which recorded its highest significant level in treatment T5 and T3, which amounted to

(60.09 and 59.84) g, respectively, which are the two treatments that were provided an addition rate of *Anemone coronaria* L. flower powder, amounting to (2) g/kg feed, with the addition of vitamin C (treatment T5) or without the addition of vitamin C (treatment T3). The egg mass rate, which achieved its highest significant levels in treatment T6, amounting to (54.83) g/bird/day. This significant improvement of the above-mentioned characteristics is due to the active substances in the powder of the flowers of the *Anemone coronaria* L. plant. The role of flavonoids is similar to that of steroid hormones [14]. These hormones work (as do flavonoids) to increase the basal metabolic rate because they are structural hormones, including estrogen, which plays an important role in promoting the growth of the oviduct and the formation of proteins necessary for the formation of the egg [15]. In turn, flavonoids stimulate the secretion of the thyroid gland and thus increase the secretion of the hormone thyroxine, which affects protein metabolism and works to increase the flow of amino acids to the ovarian cells and increase their concentration within the ovarian cells, thus working on the manufacture of proteins [16]. The results of the study also reinforce the findings of [17] that flavonoids have an important role in enhancing

the secretion of the hormone progesterone by the granulosa cells before the ovulation process. They also work in enhancing the proliferation and differentiation of the granulosa cells and improve the secretion hormones and the development of ovarian follicles. Therefore, adding it to feed is considered beneficial in enhancing the rate of Laying eggs. In other studies, the active ingredient represented by essential vegetable oils added as food supplements has been shown to significantly improve the performance of laying hens, egg production, ovary formation, and the sensory quality of eggs [18,19]. The improved performance recorded in the percentage of egg production and weight and mass of eggs in treatments containing natural antioxidants (medicinal plants) or artificial ones in this study compared to other treatments is consistent with the results of the study reached by [20].

The rate of feed consumption showed a significant difference. Treatment T5 recorded the lowest feed consumption of (93.22) g/feed per day. However, in contrast, the best feed conversion coefficient was recorded in treatments T5 and T6 (1.76 and 1.78) g of feed/g of eggs, respectively. This may be due to antioxidants, represented by the active substances found in the powder of *Anemone coronaria* L. flowers, and vitamin C alike. In line with several studies that reported an increase in egg production by up to (7%), better feed conversion by (15%) and an increase in intestinal villi by (24%) in laying hens supplemented with essential fats [21,22,23]. Essential fats are considered one of the active compounds in the flower powder used and have an antioxidant effect, enhancing immunity, egg production, and ovary formation [18], [19,24]. This may explain the low level of feed consumption for the bird to make the most of it.

This is consistent with the findings of [25] regarding the effect of using the *Gynura Procumbens* plant. The feed conversion factor for laying hens, but does not agree with the feed consumption rate.

The study also showed that there were no deaths in most of the treatments, Exception the treatment T1, in which the losses reached 20%, and treatment T3, in which the losses reached

10%. This may be due to the effective role of the food additives used as antioxidants, anti-inflammatory, anti-virus, and immune boosters [26,27]. The results of this study, which showed that the additives used improve the health of birds by reducing physical damage and mortality due to heat stress are consistent with [28], which showed that antioxidant supplements reduce the oxidative stress response in laying hens, thus supporting the hypothesis that supplemental antioxidants work. It improves chicken health by reducing the physical and physiological damage associated with heat stress.

References

- [1] Mottet, A. and Tempio, G. (2017). Global poultry production: current state and future outlook and challenges. *World's Poultry Science Journal*, 73(2): 245-256.
- [2] Huda Falih Saad, Salah Mahdi Alsudany, Sabah K.M. AL-hummod, & Azhar A. Jaffar. (2023). The impact of high temperatures on the productive performance (behavioral, physiological, and immunological) of poultry. *Texas Journal of Agriculture and Biological Sciences*, 14, 52–57.
- [3] Rojas-Downing, M. M., Nejadhashemi, A. P., Harrigan, T., Woznicki, S. A. (2017). Climate change and livestock: Impacts, adaptation, and mitigation. *Climate Risk Management*, 16, 145–163.
- [4] Emami, N.K., Jung, U., Voy, B., Dridi, S. (2020). Radical response: effects of heat stress-induced oxidative stress on lipid metabolism in the avian liver. *Antioxidants*. 10:35.
- [5] Goel, A. (2021). Heat stress management in poultry. *Journal of Animal Physiology and Animal Nutrition*, 1, 1-10.
- [6] Baojun Shi, Wei, L., Lvtong, G., Cuicui, C., Zhaonong, H., Wenjun, W. (2012). Chemical composition, antibacterial and antioxidant activity of the essential oil of *Anemone rivularis*. *Medicinal Plants Research* Vol. 6(25), pp. 4221-4224.
- [7] Mohammed S. Al Tamimi (2023). The effect of spraying with local *Anemone coronaria* flower extract and isoflurane in reducing economic losses resulting from the stress of transporting broiler chickens. PhD thesis. faculty of Agriculture. Kirkuk University.
- [8] Mohammed S. Al Tamimi; ammar Q. Shanoon; Mohsin O. Mohammed. (2023). "Effect of spraying with the extract of the local *Anemone coronaria* flowers and isoflurane in reducing the economic losses caused by stress of transporting broiler". *Kirkuk University Journal for Agricultural Sciences*, 14, 1, 41-52.

- [9] Mohammed S. Al Tamimi; ammar Q. Shanoon; Mohsin O. Mohammed. (2023). "Spraying with local anemone coronaria flower extract and isoflurane and its effect on the physiological characteristics of broiler chickens transported during different seasons". Kirkuk University Journal for Agricultural Sciences, 14, 1, 2023, 53-72.
- [10] NRC, (1994). Nutrient requirements of poultry. (9th rev. ed.). National Research Council. National Academy press, Washington, DC, USA
- [11] Duncan, D.B. (1955). Multiple ranges and multiple F-test. *Biometrics* 11: 1-42.
- [12] Ibrahim, M. I., Youssef, Klaus, M., Jürgen, Z., (2021). Effect of essential oils or saponins alone or in combination on productive performance, intestinal morphology and digestive enzymes' activity of broiler chickens. *Jan*; 105(1): 99-107.
- [13] Surai, (2014). Polyphenol compounds in the chicken/animal diet: from the past to the future. *Anim. Physiol. Anim. Nutr*, 98, pp. 19-31.
- [14] Sun, Y.X., Liu, J.C., Liu, D.Y. (2011). Phytochemicals and bioactivities of *Anemone raddeana* Regel: A review. *Pharmazie.*, 66, 813–821.
- [15] Barton, B., Herrera, G., Anamthathmakula, P., Rock, Willie, J., Harris, E., Takemaru, K., Winuthayanon, W. (2020): Roles of steroid hormones in oviductal function. *Reproduction* 159: R125-R137.
- [16] Esraa, C., Shanoon, A., AL-Dalawi, R. (2020). Effect of the use of different levels of *Gynura procumbens* leaf powder on the productive qualities of ISA BROWN white chicken. *Plant Archives* Vol. 20, Supplement 2, 2020 pp. 361-365.
- [17] Guo, Y., Li, Y., Zhang, S., Wu, X., Jiang, L., Zhao, Q., Xue, W., Huo, S., (2020). The effect of total flavonoids of *Epimedium granulosum* cell development in laying hens. *Poult. Sci.* 2020, 99, 4598–4606.
- [18] Ghajarbeygi, P., Mohammadi, A., Mahmoudi, R., Kosari-Nasab, M. (2015). *Artemisia spicigera* Essential Oil: Assessment of Phytochemical and Antioxidant Properties. *Biotechnol. Heal. Sci.* 2015;2.
- [19] Wang, H., Liang, S., Li, X., Yang, X., Long, F., Yang, X. (2019). Effects of encapsulated essential oils and organic acids on laying performance, egg quality, intestinal morphology, barrier function, and microflora count of hens during the early laying period. *Poult. Sci.* 2019; 98:6751–6760.
- [20] Amin, Kulala Wahab, Ammar Qahtan Shanoon, and Rashid Hassan Hamid Al-Dalawi (2021). The effect of adding dried Khoshnaw grape seed powder, dried Bazian tomato powder, and BHT on the productive performance of ISA BROWN laying hens. *Kirkuk University Journal for Agricultural Sciences*. Volume (12) Issue: 3. 1-14.
- [21] Cheng, H., Chen, J.F., Tang, S.G., Guo, S.C., He, C.Q., Qu, X.Y., (2022). Effects of essential oil/palygorskite composite on performance, egg quality, plasma biochemistry, oxidation status, immune response and intestinal morphology of laying hens. *Poult. Sci.* 2022, 101, 101632.
- [22] Abdel-Wareth, A.A.A., Lohakare, J.D., (2020). Productive performance, egg quality, nutrients digestibility, and physiological response of Bovans Brown hens fed various dietary inclusion levels of peppermint oil. *Anim. Feed. Sci. Technol.* 2020, 267, 114554.
- [23] Abo Ghanima, M.M., Elsadek, M.F., Taha, A.E., Abd El-Hack, M.E., Alagawany, M., Ahmed, B.M., Elshafie, M.M., El-Sabrouh, K., (2020). Effect of housing system and rosemary and cinnamon essential oils on layers' performance, egg quality, haematological traits, blood chemistry, immunity, and antioxidant. *Animals*, 10, 245.
- [24] Abou-Elkhair, R., Selim, S., Hussein, E. (2018). Effect of supplementing layer hen diet with phytochemical feed additives on laying performance, egg quality, egg lipid peroxidation and blood biochemical constituents. *Anim. Nutr.* 2018; 4:394–400.
- [25] Israa Qaiser Fares (2020). The effect of using different levels of *Gynura Procumbens* leaf powder on the productive performance, blood biochemical parameters, and intestinal flora of laying hens. Master Thesis. faculty of Agriculture. Kirkuk University.
- [26] Chambial, S., Dwivedi, S., Shukla, K. K., John, P. J., and Sharma, P. (2013). Vitamin C in disease prevention and cure: an overview. *Indian J. Clin. Biochemistr.* 28:3.
- [27] Hao, D., Xiaojie, G., Peigen, X. (2017). *Anemone* medicinal plants: Ethnopharmacology, phytochemistry and biology. *Acta Pharmaceutica Sinica* B.7(2):146–158.
- [28] Felver-Gant, J.N., Dennis, R.L., Zhao, J., Cheng, H.W. (2014). Effects of Dietary Antioxidant on Performance and Physiological Responses Following Heat Stress in Laying Hens. [Int. J. Poult Sci.](#),13(5):260-271



مقارنة تأثير إضافة مسحوق ازهار نبات *Anemone coronaria* L. المحلية وفيتامين C في العليقة على الأداء الإنتاجي للدجاج البيض المعرض للإجهاد الحراري صيفاً

عمار قحطان شعنون²

ammar.qahtan@uokirkuk.edu.iq

محمد فاضل عبد الخياط¹

mfkyatt@gmail.com

¹ وزارة البيئة، بغداد، العراق.

² قسم الإنتاج الحيواني، كلية الزراعة، جامعة كركوك، كركوك، العراق.

• البحث مستل من رسالة ماجستير للباحث الأول.

• تاريخ استلام البحث 2024/2/3 وتاريخ قبوله 2024/3/2.

الخلاصة

تم دراسة تأثير إضافة مسحوق ازهار نبات *Anemone coronaria* L. المحلية وفيتامين C على الأداء الإنتاجي للدجاج البيض المرعى في فصل الصيف والمعرض الى الاجهاد الحراري، وانجزت التجربة باستخدام (120) طائر من نوع Lohmann Brown بعمر (30) أسبوع وخلال أشهر تموز واب وأيلول ولمدة (84) يوماً حقلياً تضمنت 14 يوماً كفترة تمهيدية، قسمت التجربة الى (5) فترات إنتاجية، حيث وزعت الطيور عشوائياً الى (6) معاملات وبواقع (5) مكررات ولكل مكرر (4) طيور، اذ كانت المعاملة T1: عليقة مقارنة خالية من أي إضافة، T2: عليقة مقارنة مضاف اليها فيتامين C بنسبة 250 ملغم/كغم علف، T3: عليقة مقارنة مضاف اليها مسحوق ازهار نبات *Anemone coronaria* L. بنسبه 2 غم/كغم علف، T4: عليقة مقارنة مضاف اليها مسحوق ازهار نبات *Anemone coronaria* L. بنسبه 4 غم/كغم علف، T5: عليقة مقارنة مضاف اليها مسحوق ازهار نبات *Anemone coronaria* L. بنسبه 2 غم/كغم علف وفيتامين C بنسبة 250 ملغم/كغم علف، T6: عليقة مقارنة مضاف اليها مسحوق ازهار نبات *Anemone coronaria* L. بنسبه 4 غم/كغم علف وفيتامين C بنسبة 250 ملغم/كغم علف. اثبتت نتائج التجربة الحقلية الى وجود فروقات معنوية في المعدل العام للفترات الإنتاجية عند مستوى ($P \leq 0.05$) في جميع الصفات المدروسة، فقد سجلت المعاملة T6 اعلى نسبة انتاج بيض، واعلى معدل وزن بيض في المعاملتين T5 و T3 على التوالي، واعلى معدل كتلة بيض في المعاملة T6، واعلى معدل استهلاك علف في المعاملة T2، وأفضل معامل تحويل غذائي في المعاملة الخامسة والسادسة على التوالي، شهدت معاملة السيطرة اعلى نسبة هلاكات مع نقص في معدل الوزن الحي للطيور. حيث يلاحظ من نتائج التجربة ان اضافات مسحوق ازهار نبات *Anemone coronaria* L. وفيتامين C أدى الى التقليل من اثار الاجهاد الحراري الذي يصيب الطيور عند تعرضها الى درجات الحرارة المرتفعة صيفاً وبالتالي كان التأثير الإيجابي واضح على تحسن الاداء الإنتاجي وصحة الطيور.

الكلمات المفتاحية: شقائق النعمان الاكليلية، فيتامين C، الأداء الإنتاجي، الدجاج البيض، الاجهاد الحراري.