



The Effect of Adding Powdered Pomegranate Peels and Ginger Tubers to The Diet of Laying Hens on Some Blood Parameters

Amanj H. Mohamed ¹

akam22002@uokirkuk.edu.iq

Qana H. Al-Jabari ²

dr_qanaameen@uokirkuk.edu.iq

¹ Poultry division, Animal production department, general directorate of Garmiyah agriculture, Iraq

² Animal production department, college of agriculture, Kirkuk University, Kirkuk, Iraq.

- Date of research received 14/3/2024 and accepted 18/4/2024.

Abstract

This experiment was conducted in the fields of the Animal Production department - College of Agriculture - University of Kirkuk, from 7/24/2023 until 10/1/2023. 140 laying hens, 28 weeks old, were used. The birds were randomly distributed into 7 treatments. Each treatment included 5 replicates, with 4 birds per replicate. T1: control treatment. T2: control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed. T3: control diet with the addition of pomegranate peel powder at a concentration of 2 g/kg feed. T4: control diet with the addition of ginger tuber powder at a concentration of 1 g/kg feed. T5: control diet with the addition of ginger tuber powder at a concentration of 2 g/kg feed. T6: control diet with the addition of pomegranate peel powder at a concentration of 0.5 g/kg feed + 0.5 g/kg feed ginger tuber powder. T7: control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed + 1 g/kg feed ginger tuber powder. In the end of the experiment, blood was withdrawn from chicken for testing. The data was analyzed statistically using SAS, and the differences between the means were tested using the Duncan test at the probability level ($P < 0.05$). The present study indicates, using of 2 g/kg of pomegranate peel powder in the diet of the laying hens decrease the total protein, and Uric acid level on the blood. And also effect on the AST, MDA, and GSH level on the blood.

Keywords: Layers, Pomegranate peels, ginger tubers, powdered.

Citation: Mohammed, A., & Ameen, Q. (2024). The Effect of Adding Powdered Pomegranate Peels and Ginger Tubers to The Diet of Laying Hens on Some Blood Parameters. *Kirkuk University Journal for Agricultural Sciences*, 15(2), 47-53. doi: 10.58928/kujas.2024.147822.15205

Correspondence Author: Amanj H. Mohamed. akam22002@uokirkuk.edu.iq

Copyright: This is an open access article distributed under the terms of the creative common's attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited

Introduction:

Poultry production is one of the most pervasive industries in animal husbandry, and it is regarded as one of the fundamental and significant pillars that nations around the world rely on to ensure food security [1, 2]. Since artificial antioxidants may contain chemicals that cause cancer, there have been several attempts to use the natural dietary additives in diet of poultry [3, 4, 5, 6, and 7].

Pomegranate peels have been used because they contain natural antioxidants as well as phenolic compounds [8]. Pomegranate peels have the ability to act as antioxidants because they contain phenolic compounds and dissolved tannins in water [9]. [10] indicated that about 35% of the weight of the pomegranate fruit is made up of pomegranate peels, which are the most abundant due to their lack of use. Pomegranates have a distinguished place in folk medicine, where the sour pomegranate is used as a diuretic [11], and that the bitter taste of pomegranate peels helps in treating infections [12]. Antifungals are what distinguish ginger tubers [13], as well as containing antioxidants. It also contains phenols, alkaloids, carbohydrates, and terpenes [14].

Recent scientific research has revealed that ginger tubers possess many therapeutic properties include that it acts as antioxidants, has the ability to inhibit the formation of inflammatory compounds, and has direct anti-inflammatory effects. In addition, it is noted that ginger tubers are also effective against some types of carcinogenic diseases, regulate and control high and low blood pressure, and activate and stimulate blood circulation. It is linked to reducing heart problems and helps lower blood cholesterol [15]. The antioxidants that are naturally present in the body play a major and major role in the balance that occurs between resistance to oxidation and the production of free radicals. If there is a small amount of antioxidants that are naturally present in the body and are matched by a large amount of the free radicals produced, then what is called oxidative stress occurs, which is Its results are the destruction of genetic material, DNA, proteins, carbohydrates, and vitamins. It also

works to oxidize unsaturated fatty acids in cell membranes [16]. Antioxidants work directly or indirectly to remove the effect of, destroy, and scavenge free radicals. It is possible to classify antioxidants depending on their method of activating them.

The current study aimed to determine the effect of adding powdered pomegranate peels and ginger tubers as natural antioxidants to the diet of laying hens and their effect on some blood parameters.

Materials and methods

This experiment was conducted in the fields designated for scientific research and affiliated with the Department of Animal Production at the College of Agriculture, University of Kirkuk. It is a closed field with a solid floor containing three large-sized pullers. The experiment period was 70 days from 7/24/2023 until 10/1/2023. The rearing system was done using a vertical cage rearing system. 140 BROWN LOHMANN laying hens, 28 weeks old, were used. The birds were randomly distributed into 7 treatments. Each treatment included 5 replicates, with 4 birds per replicate. Each battery consisted of 4 floors with one feeder on each floor, installed and separated manually. Each floor consisted of five cages. Each cage contained two chickens, and the cage dimensions were (48 x 45 x 40 cm, length, width, and height, respectively) for each cage. Water was provided continuously and freely to the bird through a system of nipples connected to water supply lines, and these lines were connected to a 1000-liter water tank, and feed was provided manually at a rate of 100 g/day according to the company's guide for this breed of laying chickens. A lighting period was 16 hours per day, with a preliminary period of 15 days before starting the experiment to prepare the birds.

Powdered pomegranate peels and powdered ginger tubers were purchased from sellers of medicinal herbs in the local market of Kirkuk Governorate. The experiment diet was consist of 2708 Kcal, and 18.37 CP.

The treatments were according to below:

T1: Standard control method.

T2: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed.

T3: Standard control diet with the addition of pomegranate peel powder at a concentration of 2 g/kg feed.

T4: Standard control diet with the addition of ginger tuber powder at a concentration of 1 g/kg feed.

T5: Standard control diet with the addition of ginger tuber powder at a concentration of 2 g/kg feed.

T6: Standard control diet with the addition of pomegranate peel powder at a concentration of 0.5 g/kg feed + 0.5 g/kg feed ginger tuber powder.

T7: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed + 1 g/kg feed ginger tuber powder.

In the end of the experiment, blood was get from the hens. Fresh blood samples were withdrawn from the wing vein from the chicken and placed in test tubes devoid of anticoagulants and left for 6 hours, after which the serum was separated by a centrifuge at a speed of 3000 rpm for 15 minutes, and the serum was stored at -20°C until biochemical tests were performed.

The data was analyzed statistically using (SAS) [17], and the differences between the means were tested using the Duncan test [18] at the probability level ($P < 0.05$).

Results and discussion

The mean and standard error of total protein, total cholesterol, triglycerides, glucose, and uric acid that effected by different levels of pomegranate peel powder and ginger tubers are shown in table 1. Significant differences were found in total protein and the uric acid ($P < 0.05$). The total protein was decrease significantly in treatment three (T3) that was (4.66), and the increase in treatments 4, 6, and 7 (5.56, 5.50, and 5.74) respectively. The Current results agreed with the [19] finding when he used different levels of pomegranate peel in the diet of the laying hens. And our results disagreed with [20] who used pomegranate juice in the diet of the laying hens. And found there were no significant differences between the treatments in Total protein, total cholesterol, triglycerides, and glucose.

[21] used different levels of pomegranate peel powder in the Japanese quail diet, and they found significant differences between the treatments in each of Total protein, total cholesterol, triglycerides, and glucose.

[22] used different level of pomegranate peel on the diet of laying hen and found significant differences between the treatment on the cholesterol level, and the on the Triglycerides. Moreover [23] found there was no significant differences when he used different levels of ginger on the level of cholesterol, but the ALT, and the AS

Table (1): The effect of adding pomegranate peel powder and ginger tubers to laying hens' diet on total protein, total cholesterol, triglycerides, glucose, and uric acid.

Treatments	Blood parameters tests Birds at 40 weeks old				
	Total Protein	Total cholesterol	Triglycerides	Glucose	Uric acid
T1	5.06±0.10 ab	130.80±12.72 a	1316.50±98.21 a	206.60±3.65 a	5.80±0.73 bc
T2	5.06±0.26 ab	105.80±13.29 a	953.80±148.97 a	222.80±10.57 a	5.80±1.68 bc
T3	4.66±0.16 b	99.40±13.14 a	939.20±154.30 a	217.40±7.78 a	4.40±0.40 c
T4	5.56±0.17 a	117.80±20.95 a	1111.40±179.73 a	215.40±12.26 a	7.40±1.17 b
T5	5.06±0.30 ab	107.00±21.74 a	1189.80±119.40 a	201.20±8.63 a	7.40±1.94 b
T6	5.50±0.18 a	119.60±20.94 a	1350.80±69.88 a	230.00±6.42 a	10.80±1.36 a
T7	5.74±0.26 a	131.40±21.16 a	1039.00±154.26 a	277.20±62.47 a	9.80±1.93 ab
Level of Sig.	*	NS	NS	NS	*

T1: Standard control method. T2: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed. T3: Standard control diet with the addition of pomegranate peel powder at a concentration of 2 g/kg feed. T4: Standard control diet with the addition of ginger tuber powder at a concentration of 1 g/kg feed. T5: Standard control diet with the addition of ginger tuber powder at a concentration of 2 g/kg feed. T6: Standard control diet with the addition of pomegranate peel powder at a concentration of 0.5 g/kg feed + 0.5 g/kg feed ginger tuber powder. T7: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed + 1 g/kg feed ginger tuber powder.

The mean and standard error of the liver enzymes (AST, and ALT) antioxidants (MDA, GSH) that effected by different levels of pomegranate peel powder and ginger tubers are shown in table 2. Significant differences were found between the treatments in AST, MDA, and GSH. The AST was decrease significantly in treatment three (T3) that was (156.00), and was increase in treatment five (T5). The MDA was decrease significantly in treatment one, and two (2.05, and 2.06) respectively, and increase in both treatment 6, and 7 (3.21, and 3.45) respectively. The GSH was increase in T1

(4.15), and decrease significantly in all the other treatments. The Current results agreed with the [24] finding when he used different levels of pomegranate peel in the diet of the laying hens. [25] used the oil seed of pomegranate, and he found significant differences among the treatments in GSH of the laying hens. [22] used different level of pomegranate peel on the diet of laying hen and found significant differences between the treatments on the GSH. Moreover [26] found there was significant differences when he used different levels of ginger on the level of ALT, AST and the MDA.

Table (2): The effect of adding pomegranate peel powder and ginger tubers to laying hens' diet on liver enzymes (AST, ALT) and antioxidants (MDA, GSH).

Treatments	Blood parameters tests			
	Birds at 40 weeks old			
	AST U/L	ALT U/L	MDA $\mu\text{mol/L}$	GSH $\mu\text{mol/L}$
T1	168.80 \pm 8.72 bc	2.82 \pm 0.50 a	2.05 \pm 0.14 b	4.15 \pm 0.26 a
T2	185.20 \pm 13.68 ab	0.84 \pm 0.30 a	2.06 \pm 0.17 b	2.44 \pm 0.25 b
T3	156.00 \pm 35.09 c	1.74 \pm 0.81 a	2.95 \pm 0.48 ab	1.70 \pm 0.37 b
T4	175.80 \pm 3.62 b	2.40 \pm 0.99 a	2.64 \pm 0.45 ab	2.15 \pm 0.88 b
T5	223.00 \pm 7.60 a	2.54 \pm 2.18 a	3.06 \pm 0.37 ab	1.34 \pm 0.29 b
T6	211.60 \pm 12.10 ab	3.24 \pm 1.56 a	3.21 \pm 0.21 a	1.44 \pm 0.36 b
T7	189.20 \pm 6.90 ab	1.42 \pm 0.60 a	3.45 \pm 0.33 a	1.35 \pm 0.32 b
Level of Sig.	*	NS	*	*

T1: Standard control method. T2: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed. T3: Standard control diet with the addition of pomegranate peel powder at a concentration of 2 g/kg feed. T4: Standard control diet with the addition of ginger tuber powder at a concentration of 1 g/kg feed. T5: Standard control diet with the addition of ginger tuber powder at a concentration of 2 g/kg feed. T6: Standard control diet with the addition of pomegranate peel powder at a concentration of 0.5 g/kg feed + 0.5 g/kg feed ginger tuber powder. T7: Standard control diet with the addition of pomegranate peel powder at a concentration of 1 g/kg feed + 1 g/kg feed ginger tuber powder.

Conclusion

The present study indicates, using of 2 g/kg of pomegranate peel powder in the diet of the laying hens decrease the total protein, and Uric acid level on the blood. And also effect on the AST, MDA, and GSH level on the blood.

References

- [1] Al-Jabari, Q. H., & Shaker, A. S. (2023). The Effect of Adding Moringa Leaf Powder to the Adapted Quail Diet During the Egg Production Stage on the Productive Performance and some Biochemical Blood Characteristics. In IOP Conference Series: Earth and Environmental Science, 1262(7): 072052
- [2] Hassan, A. O. , Al- Jabari, Q. H. A. , Mustafa, N. A. (2023). Impact of Adding Chitosan and Probiotic to Broiler Dietary on carcass traits. Kirkuk University Journal For Agricultural Sciences, 14(3):75-81.
- [3] AL-Khaldani, C., & Ameen, Q. (2022). Effect of fodder addition of Moringa oleifera leaf powder and probiotic on the productive characteristics of broilers. *Kirkuk University Journal For Agricultural Sciences*, 13(3), 49-61.
- [4] AL-Khaldani, C., & Ameen, Q. (2022). Effect of adding Moringa oleifera leaf powder with or without probiotic on growth performance, carcass characteristics and some biochemical blood characteristics for broiler. *Kirkuk University Journal For Agricultural Sciences*, 13(3), 186-201.
- [5] Al-Qaisi, A. H. M. , Al-Jabari, Q. H. A. (2023).The effect of partial or total substitution of raw or roasted domestic sesame seeds in laying hens on the qualitative qualities of eggs. *Kirkuk University Journal For Agricultural Sciences*. 14(3):113-122.
- [6] AL-Jabari, Q. H., Mohammed, M. A., Baker, A. G., & Shaker, A. S. (2024). Effect of Supplementation of Different Levels of Azolla Plant Powder on The Productive Performance and Some Qualitative Characteristics of Eggs of Japanese Quail. *Egyptian Journal of Veterinary Sciences*, 55(7), 1869-1874.
- [7] AL-Jabari, Q. H. H., Baker, A. G., Amen, S. H., & Shaker, A. S. (2024). The Effect of Adding Azolla Plant Powder in Quail Diets on The Carcass Traits and Some Blood Traits. *Egyptian Journal of Veterinary Sciences*, 55(5), 1251-1255.
- [8] Mo, Y., Ma, J., Gao, W., Zhang, L., Li, J., Li, J., Zang, J. (2022). Pomegranate Peel as a Source of Bioactive Compounds: A Mini Review on Their Physiological Functions. *Front Nutr*. 9:887113.
- [9] Altunkaya-Dinçay, Arzu. (2014). Effect of Grape Leaf Extract on Phenolic Profile and Browning of Fresh-Cut Lettuce (*Lactuca sativa*). *Journal of Food Processing & Preservation*. 38.
- [10] Elfalleh, W., Tlili, N., Nasri, N., Yahia, Y., Hannachi, H., Chaira, N., Ying, M., Ferchichi, A. (2011). Antioxidant Capacities of Phenolic Compounds and Tocopherols from Tunisian Pomegranate (*Punica granatum*) Fruits. *Journal of food science*. 76.
- [11] Mohammadi, M., Boghrati, Z., Emami, S.A., Akaberi, M. (2023). Pomegranate: A review of the

heavenly healer's past, present, and future. Iran J Basic Med Sci. 26(11):1245-1264.

[12] El-Beltagi, H.S., Eshak, N.S., Mohamed, H.I., Bendary, E.S.A., Danial, A.W. (2022) Physical Characteristics, Mineral Content, and Antioxidant and Antibacterial Activities of Punica granatum or Citrus sinensis Peel Extracts and Their Applications to Improve Cake Quality. Plants (Basel). 11(13):1740.

[13] Toubia, E. P., Zakaria, M., and Tahereh, E. (2012). Anti-fungal activity of cold and hot water extracts of spices against fungal pathogens of Roselle (*Hibiscus sabdariffa*) in vitro. Microbial Pathogenesis, 52(2), 125-129.

[14] Grzanna, R., Lindmark, L., & Frondoza, C. G. (2005). Ginger--an herbal medicinal product with broad anti-inflammatory actions. Journal of medicinal food, 8(2), 125-132.

[15] Fakhri, S., Patra, J. K., Das, S. K., Das, G., Majnooni, M. B., & Farzaei, M. H. (2021). Ginger and Heart Health: From Mechanisms to Therapeutics. Current molecular pharmacology, 14(6), 943-959.

[16] Gholami-Ahangaran, M., Karimi-Dehkordi, M., Akbari Javar, A., Haj Salehi, M., Ostadpoor, M. (2021). A systematic review on the effect of Ginger (*Zingiber officinale*) on improvement of biological and fertility indices of sperm in laboratory animals, poultry and humans. Vet Med Sci. 7(5):1959-1969.

[17] SAS. (2005). SAS/STAT' User's Guide for Personal Computers. Release 8.2. SAS Institute Inc. Cary, NC, USA.

[18] Duncan, D. B. (1955). Multiple range and multiple F test. Biometrics, 11: 1-42.

[19] Jabber, S. S., Khalil, A. H. and Al-Gharawi, J. K. (2021). Effect of Adding Different Levels of Pomegranate Peel Powder to Feed on Some Biochemical Blood Traits of Laying Hens. IOP Conf. Series: Earth and Environmental Science 923: 012034.

[20] GULTEPE, E. E., IQBAL, A., ÇETİNGÜL, İ. S., UYARLAR, C., et al. (2022). Effects of pomegranate (*Punica granatum L.*) juice as a short-term water supplement during the peak production cycle in laying hens. Ankara Üniversitesi Veteriner Fakültesi Dergisi, 69(3), 241-249.

[21] Abbas R. J., Al-Salhie, K. Ch. K., and Al-Hummod, S. K. M. (2017). The effect of using different levels of pomegranate (*Punica granatum*) peel powder on productive and physiological performance of Japanese quail (*Coturnix coturnix japonica*). *Livestock Research for Rural Development. Volume 29, Article #231.*

[22] Eid, Y., Kirrella, A.A., Tolba, A., El-Deeb, M., Sayed, S., El-Sawy, H.B., Shukry, M., and Dawood M.A.O. (2021). Dietary Pomegranate By-Product Alleviated the Oxidative Stress Induced by Dexamethasone in Laying Hens in the Pre-Peak Period. Animals (Basel). 11(4):1022.

[23] Wen, C., Y. Gu. Z. Ta, Z. Cheng, T. Wang, and Y. Zhou. (2019). Effects of Ginger Extract on Laying Performance, Egg Quality, and Antioxidant Status of Laying Hens. Animals, 9: 857

[24] Xu, P., Wang, J., Chen, P., Ding, H., Wang, X., Li, S., Fan, X., Zhou, Z., Shi, D., Li, Z., Cao, S., and Xiao, Y. (2024). Effects of pomegranate (*Punica granatum L.*) peel on the growth performance and intestinal microbiota of broilers challenged with *Escherichia coli*. Poultry science, 103(2), 103304.

[25] Bölükbaşı, Ş.C., Dumlu, B., Yağanoğlu, A.M., (2023). Improved biological value of eggs due to the addition of pomegranate seed oil to laying-hen diets. Arch Anim Breed. 66(1):121-129.

[26] Abd El-Hack ME, Alagawany M, Shaheen H, Samak D, Othman SI, Allam AA, Taha AE, Khafaga AF, Arif M, Osman A, El Sheikh AI, Elnesr SS, Sitohy M. Ginger and Its Derivatives as Promising Alternatives to Antibiotics in Poultry Feed. Animals (Basel). 2020 Mar 9;10(3):452.



تأثير إضافة مسحوق قشور الرمان ودرنات الزنجبيل إلى علائق الدجاج البياض في بعض صفات الدم

قانع حسين امين الجباري²

dr_qanaameen@uokirkuk.edu.iq

أمانج هلال محمد¹

akam22002@uokirkuk.edu.iq

¹ شعبة الدواجن، قسم الإنتاج الحيواني، المديرية العامة للزراعة كرميان، العراق.

² قسم الإنتاج الحيواني، كلية الزراعة، جامعة كركوك، كركوك، العراق.

• تاريخ استلام البحث 2024/3/14 وتاريخ قبوله 2024/4/18.

الخلاصة

أجريت هذه التجربة في الحقول المخصصة للبحث العلمي والتابعة لقسم الإنتاج الحيواني في كلية الزراعة جامعة كركوك. مدة التجربة 70 يوماً من 2023/7/24 حتى 2023/10/1. تم استخدام 140 دجاجة بياضة من نوع براون لوهمان بعمر 28 اسبوع. وزعت الطيور عشوائياً على 7 معاملات. تضمنت كل معاملة 5 مكررات، بواقع 4 طيور في كل مكرر. المعاملات المستخدمة كانت كالاتي: المعاملة الاولى: معاملة السيطرة، المعاملة الثانية: إضافة مسحوق قشر الرمان بتركيز 1 غرام/كغم علف، المعاملة الثالثة: إضافة مسحوق قشر الرمان بتركيز 2 غرام/كغم علف، المعاملة الرابعة: إضافة مسحوق درنات الزنجبيل بتركيز 1 غرام/كغم علف، المعاملة الخامسة: إضافة مسحوق درنات الزنجبيل بتركيز 2 غرام/كغم علف، المعاملة السادسة: إضافة مسحوق قشر الرمان بتركيز 0.5 غرام/كغم علف + 0.5 غرام/كغم علف مسحوق درنات الزنجبيل، المعاملة السابعة: إضافة مسحوق قشر الرمان بتركيز 1 غرام/كغم علف + 1 غرام/كغم علف مسحوق درنات الزنجبيل. وفي نهاية التجربة، تم سحب الدم من الدجاج و تخزين المصل عند -20 درجة مئوية حتى إجراء الاختبارات البيوكيميائية. وتم تحليل البيانات إحصائياً باستخدام SAS، وتم اختبار الفروق بين المتوسطات باستخدام اختبار دنكان عند مستوى الاحتمال $P < 0.05$. أشارت الدراسة الحالية إلى أن استخدام 2 غرام/كغم من مسحوق قشر الرمان في عليقة الدجاج البياض قد ادى الى خفض مستوى البروتين الكلي وحمض البوليك في الدم. وأيضاً التأثير على مستوى AST وMDA وGSH في الدم.

الكلمات المفتاحية: الدجاج البياض، قشور الرمان، درنات لزنجبيل، مسحوق.