



Impact of Different Zinc Levels and Benzyl Adenine Concentrations Applied as Foliar Spray on the Vegetative Growth of Two *Zinnia elegans* Cultivars.

Layla Shaaban Mohammed¹

Ban Jibrael Khoshaba¹

Niroosh Sulaiman Hasan¹

Amira abd el-hamid Yousef²

¹Department of Horticulture, College of Agriculture Engineering Sciences, University of Duhok, Kurdistan region, IRAQ.

²Department of Botanical Gardens, Agricultural Research Center, Egypt.

*Corresponding Author: layla.mohammed@uod.ac.

Received:27/05/2025

Revised: 19/06/2025

Accepted: 05/08/2025

Published: 01/09/2025

ABSTRACT

This study was conducted at the Department of Horticulture, College of Agriculture Engineering Sciences. To investigate the effects of varying Zn concentrations (0, 0.50%, and 0.75%), as well as varying BA concentrations (0.50, and 100) mg⁻¹ on some vegetative traits of two cultivars of Zinnia plant. The results showed that: The mean values showed that the largest plants (90.556 cm) were recorded in the *Zinnia Lavonda* cultivar. Also, the same cultivar shows the highest number of branches, fresh and dry aerial weight, and total carotene reached (8.160, 213.194 g, 11.534 g, and 12.398g), respectively. While another parameter such as leaves number, internodes number, leaf area, and total chlorophyll were observed in the *Zinnia rosso scarlatta* cultivar and gave the maximum results reached (23.28, 10.79, 19.20, and 30.56), respectively. Also, data reported in the same table reveal that leaves number, branches number, number of internodes, shoot fresh and dry weight, leaf area (cm²), and finally total carotene increased significantly with a 50 mg⁻¹ concentration of BA for the *Zinnia Lavanda* cultivar (26.741, 9.296, 12.556, 259.054, 13.829, 24.489, and 14.798 cm²), respectively, compared to the control. About the three factors, the highest plant high for two cultivars *Zinnia Lavanda* and *Zinnia rosso scarlatta*, reached 101.411 cm when spraying the plant by 50 mg⁻¹ + 0.50% Zn and 96.622 cm at 0.50% Zn alone, respectively, than the control plants (77.856 and 77.356 cm), respectively, were obtained in two cultivars. While the number of leaves, number of branches, number of internodes, and leaf area (cm²) increased significantly compared to control and other treatments, the main value reached (30.889, 9.778, 15.222, and 33.230 cm²) respectively, when the *Zinnia Lavanda* cultivar was spraying with 100 mg⁻¹BA+ 0.50% Zn. Also, the plant fresh weight, plant dry weight, total chlorophyll, and total carotene give the height value reached (333.080g, 15.680g, 42.714 mg.g⁻¹, and 17.90 mg.g⁻¹), respectively, when the *Zinnia Lavanda* cultivar was spraying with 100 mg⁻¹BA+ 0.75% Zn.

Keywords: Levels of Zn, BA Concentrations, Spraying, Vegetative Growth and Two Cultivars.

Copyright © 2025. This is an open-access article distributed under the Creative Commons Attribution License.

INTRODUCTION

Zinnia elegans is native to Mexico and Central America and is a member of the Asteraceae family. Due to its ability to tolerate hot, dry weather, it is widely planted for commercial purposes as a cut flower and bedding plant [1]. *Zinnia* is a crop with economic value as well. *Zinnia* may be studied to determine its potential as a dry condition cut flower crop because of its natural ability to survive in both dry and humid conditions. Due to the variety of colors and long flowering period, *Zinnia elegans* is a beautiful ornamental plant that may be utilized in landscape design. [2]. It grows after the frost season ends and is susceptible to low temperatures. *Z. elegans* are very easy to propagate by seeds, and it is used as a cut flower [3]. It's one of those plants that's frequently utilized to create flower arrangements in urban landscaping. *Zinnias* look great when planted alone or in combination with other ornamental annuals [4, 5]. Furthermore, *Zinnia elegans* is also suitable for cutting to make bouquets arrangements and for growing in pots and containers [6, 7]. This plant requires minimal care and blends in perfectly with the current trend of natural gardens. *Zinnia elegans* is a heliophyte that tolerates low air humidity as well as high soil temperatures. *Zinnia* is a usual long-day plant. [8, 9, 10, 11]. *Zinnia* is the two most promising between 20-30 species [6], *Zinnia elegans* is the parent plant of many beautiful cultivars, including a wide range of types that are commonly used for their attractive flowers. It

grows easily in rock gardens, containers for windows, and beds [12]. Providing an appropriate amount of nutrition during the growth season is one of the most crucial aspects of producing cut flowers [13]. Flowering crops' vase life can be effectively extended by using plant growth regulators, several of which have been shown to increase flower vase life. [14]

Small quantities of plant growth regulators can modify physiological processes in plants and are essential for elongation and flower formation, among other aspects of plant growth and development. [15]. When using plant growth regulators, controlling blooming is one of the most crucial practical issues. Plant growth regulators play from seed germination until the stage of senescence [16]. Cytokinin can be utilized in different applications, such as the treatment of seeds [17] to applications during flowering [18]. Synthetic cytokinin like benzyl adenine (BA) may improve plant development by promoting cell division, breaking dormancy in the buds, and improving the lateral bud development [19]. When synthetic cytokinin was applied to leaves, it enhanced the production of proteins and photosynthesis pigment in the tissues. [20].

Zinc (Zn) is a crucial micronutrient for higher plants, as it is needed for the work of various enzymes (like dehydrogenases and RNA and DNA polymerases), gene expression, the health of bio-membranes, carbon metabolism in photosynthesis, carbohydrate metabolism, and protein production. Additionally, zinc is more important for biomass generation; a zinc deficit may impact on thylakoids' photochemical functions, preventing the biophysical processes of photosynthesis. [21]. In the case of excess Zn, one way to reduce damage is to prevent the uncontrolled oxidation caused by antioxidant enzymes, which require Fe, Mn, Cu, and Zn as cofactors of metals. With the introduction of micronutrients that are necessary for plant growth, it became achievable to obtain the maximum amount of plant growth [22]. Micronutrients greatly impacted plant growth and development, including zinc nutrients. Zinc has a beneficial effect on many ornamental plants and is essential for plant physiology, where it stimulates certain enzymes involved in the metabolism of carbohydrates. [23]. Zinc sprayed on the foliage increased by 0.5% and had the greatest impact on the size of the increase and vegetative growth [24].

The aim of an experiment was to enhance the growth of two cultivars of Zinnia. To investigate the impact of concentrations of synthetic cytokinin BA and Zinc levels alone and their interactions with BA on the growth performance of the two cultivars of the Zinnia plant.

MATERIALS AND METHOD

The study was conducted from April 2024 at the Department of Horticulture's Lath House, College of Agriculture Engineering Sciences, University of Duhok, Kurdistan region, Iraq. The study was arranged in a factorial test with a RCBD design with three factors and three replicates, each replicate consisting of four plants. The different materials used include BA (0, 50, and 100 mg l⁻¹) with ZnSO₄ (0, 0.50, and 0.75%), % used alone, and their interaction on the growth of two cultivars of the Zinnia plant (Lavanda and rosso scarlatta). So, the experiment includes (3×3×2×3×4=216) plants. The Pagano Costantino company in Italy supplied the Zinnia (*Zinnia elegans*) seeds, which were imported via Kurdistan offices, one of the Duhok's agricultural offices. At the beginning the seeds were sowing in peatmoss. After germination, the seedlings were planted in 24-cm containers in a growth medium that contained peatmoss and river soil at a 2:1 v/v ratio. The pots were placed in a lather with an average day/night temperature of 25°C/18°C. The plants at the four-leaf stage were sprayed individually with different concentrations of this BA and Zn, which are used in foliar application, by spraying the plant three times and 15 days between the sprays. Study parameters: The study assessed various growth parameters, including plant height (cm), leaves number per plant, branches number per plant, internodes number per plant, shoot fresh weight, shoot dry weight, and leaf area (cm²), total chlorophyll (mg/100g⁻¹ fresh weight) and total carotene (mg/100g⁻¹ fresh weight).

Statistical Analysis

An RCBD design was used to conduct the experiment. Three replications of each treatment were used, with four plants in each replication. Analysis of variance (ANOVA) was performed on the collected data, and the Duncan test was used to determine the mean values at P < 0.05 using the SAS program.

Results

Effect of cultivars

The results in Table 1 regarding many vegetative parameters such as plant height (cm), number of leaves. Plant⁻¹, number of branches. Plant⁻¹, number of internodes. Plant⁻¹, aerial fresh and dry weight, leaf area (cm²), and finally total chlorophyll, total carotene showed that the effect of zinnia cultivars was non-significant in all traits except the number of branches; it was significant between the cultivars.

Table (1): Effect of different cultivars on growth characteristics of zinnia plants

Cultivar	plant height (cm)	leaves number. Plant ⁻¹	branches number. Plant ⁻¹	internodes Number. Plant ⁻¹	shoot fresh weight	shoot dry weight	leaf area (cm ²)	total chlorophyll (mg/100g ⁻¹ fresh weight)	total carotene (mg/100g ⁻¹ fresh weight)
Zinnia Lavonda	90.556 ^a	22.469 ^a	8.160 ^a	10.691 ^a	213.194 ^a	11.534 ^a	19.177 ^a	25.754 ^b	12.398 ^a
Zinnia rosso scarlatta	90.464 ^a	23.284 ^a	7.185 ^b	10.790 ^a	181.890 ^a	10.075 ^a	19.206 ^a	30.561 ^a	12.093 ^a

The mean values showed that the largest plants (90.556 cm) were recorded in the Zinnia Lavonda cultivar. Also, the same cultivar shows the maximum number of branches, aerial fresh and dry weight, and total carotene reached (8.160, 213.194 g, 11.534 g, and 12.398), respectively. While another parameter such as number of leaves, number of internodes, leaf area, and total chlorophyll were observed in the Zinnia rosso scarlatta cultivar and gave the maximum results reached (23.284, 10.790, 19.206, and 30.561), respectively.

Effect of BA concentration on different characteristics of two cultivars of zinnia plants

The data illustrated in Table 2 indicates that rising the BA concentrations significantly increased zinnia parameters such as plant height (cm) and leaves number. Plant⁻¹, branches number. Plant⁻¹, internodes number. Plant⁻¹, shoot fresh and dry weight, leaf area (cm²), and finally total chlorophyll, total carotin compared to untreated plants. The tallest plants obtained from the treatment of 50 mg⁻¹ BA reached 96.556 cm for the Zinnia Lavanda, followed by Zinnia rosso-scarlatta cultivars, whose maximum length reached 25.778 cm at 100 mg⁻¹ BA. About the other characters, data reported in the same table reveal that number of leave, number of branches, number of internodes, plant fresh and dry weight, leaf area (cm²), finally total carotin increased significantly with a 50 mg⁻¹ concentration of BA for the Zinnia Lavanda cultivar (26.741, 9.296, 12.556, 259.054, 13.829, 24.489, and 14.798), respectively, compared to the control. Furthermore, the highest value of number of leaves. Plant⁻¹, number of branches. Plant⁻¹, number of internodes. Plant⁻¹, plant fresh and dry weight, leaf area (cm²), finally total carotin was recorded for the Zinnia rosso scarlatta cultivar at 100 mg⁻¹ BA with a significant increase compared to the control (25.778, 7.852, 12.370, 203.083, 12.833, 22.930, and 13.863), respectively, compared to the control. Also, 50 mg⁻¹ BA spraying in the Zinnia rosso scarlatta cultivar gave the highest value of total chlorophyll with significant differences compared to the other treatment, and the control reached 34.789.

Table (2): Effect of foliar application of BA concentration on different characteristics of two cultivars of zinnia plants

Cultivar	BA mg ⁻¹	plant height (cm)	leaves number. Plant ⁻¹	branches number. Plant ⁻¹	internodes Number. Plant ⁻¹	shoot fresh weight	shoot dry weight	leaf area (cm ²)	total chlorophyll (mg/100g ⁻¹ fresh weight)	total carotene (mg/100g ⁻¹ fresh weight)
Zinnia Lavanda	0	85.126 ^c	17.185 ^b	6.704 ^c	8.074 ^c	130.661 ^b	7.342 ^b	10.151 ^c	11.901 ^c	8.650 ^b
	50	96.556 ^a	23.481 ^a	8.481 ^{ab}	11.444 ^{ab}	249.868 ^a	13.432 ^a	22.892 ^a	32.374 ^a	13.746 ^a
	100	89.985 ^{abc}	26.741 ^a	9.296 ^a	12.556 ^a	259.054 ^a	13.829 ^a	24.489 ^a	32.987 ^a	14.798 ^a
Zinnia rosso scarlatta	0	88.141 ^{bc}	21.926 ^{ab}	6.444 ^c	9.222 ^{bc}	127.933 ^b	6.689 ^b	16.144 ^b	26.342 ^b	9.564 ^b
	50	89.481 ^{bc}	22.148 ^{ab}	7.259 ^{bc}	10.778 ^{abc}	214.654 ^a	10.702 ^{ab}	18.543 ^b	34.789 ^a	12.851 ^a
	100	93.769 ^{ab}	25.778 ^a	7.852 ^{abc}	12.370 ^a	203.083 ^a _b	12.833 ^a	22.930 ^a	30.551 ^{ab}	13.863 ^a

Effect of Zinc level on different characteristics of two cultivars of zinnia plants

The data regarding some growth parameters are shown in Table 2. The results showed that such as plant height

(cm) and number of leaves. Plant⁻¹, number of branches. Plant⁻¹, number of internodes. Plant⁻¹, plant fresh and dry weight, leaf area (cm²), and finally total chlorophyll and total carotin were significantly affected by zinc application. The results indicated that Zn levels significantly varied the whole parameter of two *Zinnia* cultivars compared to the control. The main values of the data showed greater plant height and number of leaves. Plant⁻¹, number of branches., number of internodes, plant fresh and dry weight, leaf area (cm²), and finally total chlorophyll, total carotin (94.837, 26.444, 8.333, 12.481, 262.440, 12.149, 24.817, 32.20, and 14.562), respectively, were recorded for *Zinnia Lavanda* cultivar with application of Zn at the rate 50% and (94.156, 26.889, 7.741, 12.593, 244.617, 11.908, 26.134, 39.749, and 14.145) respectively, were recorded for *Zinnia rosso scarlatta* cultivar with application of Zn at the rate 100%.

Table (3): Effect of foliar application of Zn % on different characteristics of two cultivars of *Zinnia* plants

Cultivar	Zn%	plant height (cm)	leaves number. Plant ⁻¹	branches number. Plant ⁻¹	internodes Number. Plant ⁻¹	shoot fresh weight	shoot dry weight	leaf area (cm ²)	total Chlorophyll (mg/100g ⁻¹ fresh weight)	total Carotene (mg/100g ⁻¹ fresh weight)
Zinnia Lavanda	0.00	84.470 ^c	16.296 ^b	7.519 ^{ab}	7.926 ^b	143.869 ^c	9.372 ^{ab}	8.491 ^b	13.147 ^c	8.527 ^b
	0.50	94.837 ^a	26.444 ^a	8.333 ^a	12.481 ^a	262.440 ^a	12.149 ^a	24.817 ^a	32.208 ^b	14.562 ^a
	0.75	92.359 ^{ab}	24.667 ^a	8.630 ^a	11.667 ^a	233.274 ^{ab}	13.082 ^a	24.224 ^a	31.907 ^b	14.105 ^a
Zinnia rosso scarlatta	0.00	86.177 ^{bc}	16.741 ^b	6.630 ^a	8.296 ^b	141.797 ^c	6.876 ^b	9.214 ^b	14.812 ^c	9.652 ^b
	0.50	91.059 ^{abc}	26.222 ^a	7.185 ^{ab}	11.481 ^a	159.258 ^{bc}	11.441 ^{ab}	22.269 ^a	37.122 ^a	12.482 ^a
	0.75	94.156 ^a	26.889 ^a	7.741 ^{ab}	12.593 ^a	244.617 ^a	11.908 ^a	26.134 ^a	39.749 ^a	14.145 ^a

Effect of BA concentration and Zn% on different characteristics of two cultivars of zinnia plants

The variability of growth parameters of zinnia cultivars in response to BA concentrations and Zn % foliar application is shown in table 4. In terms of the maximum plant height for two cultivars of (*Zinnia Lavanda* and *Zinnia rosso scarlatta*), the higher values reached 101.411 cm when spraying the plant by 50 mg l⁻¹ BA+0.50 % Zn and 96.622 cm at 0.50 % Zn alone respectively than the control plants (77.856 and 77.356 cm) respectively were obtained in two cultivars.

Table (4): Effect of foliar application of BA concentrations and Zn% on different characteristics of two cultivars of zinnia plants

Cultivar	BA mg l ⁻¹	Zn%	plant height (cm)	leaves number. Plant ⁻¹	branches number. Plant ⁻¹	internodes Number. Plant ⁻¹	leaf area (cm ²)
Zinnia Lavanda	0	0.00	77.856 ^{de}	12.667 ^e	5.333 ^f	6.111 ^d	8.735 ^e
		0.50	90.311 ^{a-d}	19.333 ^{cde}	6.889 ^{b-f}	9.667 ^{a-d}	9.555 ^e
		0.75	87.211 ^{b-e}	19.556 ^{b-e}	7.889 ^{a-f}	8.444 ^{cd}	12.163 ^{de}
	50	0.00	89.711 ^{a-e}	17.111 ^{de}	8.667 ^{a-d}	8.889 ^{bcd}	8.930 ^e
		0.50	101.411 ^a	29.111 ^{abc}	8.333 ^{a-e}	12.556 ^{abc}	31.667 ^a
		0.75	98.544 ^{ab}	24.222 ^{a-d}	8.444 ^{a-e}	12.889 ^{abc}	28.080 ^{ab}
	100	0.00	85.844 ^{b-e}	19.111 ^{cde}	8.556 ^{a-d}	8.778 ^{bcd}	7.808 ^e
		0.50	92.789 ^{abc}	30.889 ^a	9.778 ^a	15.222 ^a	33.230 ^a
		0.75	91.322 ^{abc}	30.222 ^a	9.556 ^{ab}	13.667 ^{abc}	32.429 ^a
Zinnia rosso scarlatta	0	0.00	77.356 ^e	17.111 ^{de}	5.667	7.778 ^{bcd}	8.433 ^e
		0.50	96.622 ^{abc}	22.889 ^{a-d}	7.000 ^{a-f}	8.889 ^{bcd}	18.023 ^{cd}
		0.75	90.444 ^{a-d}	25.778 ^{a-d}	6.667 ^{c-f}	11.000 ^{a-d}	21.975 ^{bc}
	50	0.00	85.256 ^{cde}	16.000 ^{de}	7.889 ^{a-f}	8.111 ^a	9.574 ^e
		0.50	87.167 ^{b-e}	25.111 ^{a-d}	6.778 ^{a-f}	12.000 ^{a-d}	19.319 ^c
		0.75	96.022 ^{abc}	25.333 ^{a-d}	7.111 ^{a-f}	12.222 ^{abc}	26.737 ^{ab}
	100	0.00	95.920 ^{abc}	17.111 ^{de}	6.333 ^{def}	9.000 ^{bcd}	9.635 ^e
		0.50	89.389 ^{a-e}	30.667 ^a	7.778 ^{a-f}	13.556 ^{abc}	29.464 ^a

0.75	96.000 ^{abc}	29.556 ^{ab}	9.444 ^{abc}	14.556 ^{ab}	29.690 ^a
------	-----------------------	----------------------	----------------------	----------------------	---------------------

While number of leaves, number of branches, number of internodes and leaf area (cm^2) increased significantly compared to control and other treatment, the main value reached (30.889, 9.778, 15.222 and 33.230 cm^2) respectively when the Zinnia Lavanda cultivar spraying with 100 mg l^{-1} BA+ 0.50% Zn. Concerning the combination of the three factors BA concentration and Zn% on different characteristics of Zinnia rosso scarlatta it was demonstrated that the Zinnia rosso scarlatta cultivar under spraying in 100 mg l^{-1} BA and 0.75 % Zn, which recorded the maximum value for number of branches, number of internodes, (9.444, 14.556) respectively. However, the maximum number of leaves 30.667 was noticed at 100 mg l^{-1} BA and 0.50 % Zn application.

Also, BA concentrations and Zn % foliar application are shown in table 5. the plant fresh weight, Plant dry weight, total chlorophyll and total carotin give the height value reached (333.080, 15.680, 42.714 and 17.90) respectively, when the Zinnia Lavanda cultivar sprayed with 100 mg l^{-1} BA+ 0.75% Zn.

Concerning the combination of the three factors BA concentration and Zn% on different characteristics of Zinnia rosso scarlatta it was demonstrated that the Zinnia rosso scarlatta cultivar under spraying in 100 mg l^{-1} BA and 0.75 % Zn, which recorded the maximum value for plant fresh weight, leaf area (cm^2) and total carotin reached (342.507, 29.690 and 17.112) respectively. However, the maximum plant dry weight (15.003g) was noticed at 100 mg l^{-1} BA and 0.50 % Zn application, The total chlorophyll generally increased, but 50 mg l^{-1} BA and 0.50 % Zn concentrations had no statistically significant effect on the total carotin

Table (5): Effect of foliar application of BA concentrations and Zn% on different characteristics of two cultivars of zinnia plants

Cultivar	BA mg l^{-1}	Zn%	shoot fresh weight	shoot dry weight	total chlorophyll (mg/100g^{-1} fresh weight)	total carotene (mg/100g^{-1} fresh weight)
Zinnia Lavanda	0	0.00	113.620 ^c	6.253 ^{ab}	8.689 ^d	7.486
		0.50	137.233 ^c	7.793 ^{ab}	13.506 ^d	9.539
		0.75	141.130 ^c	7.980 ^{ab}	13.508 ^d	8.926
		0.00	176.653 ^{bc}	10.430 ^{ab}	15.736 ^d	9.441
	50	0.50	347.337 ^a	14.280 ^{ab}	41.886 ^{ab}	16.317 ^{ab}
		0.75	225.613 ^{abc}	15.587 ^a	39.499 ^{abc}	15.482 ^{ab}
	100	0.00	141.333 ^c	11.433 ^{ab}	15.015 ^d	8.654 ^c
		0.50	302.750 ^{ab}	14.373 ^a	41.231 ^{abc}	17.831 ^a
		0.75	333.080 ^a	15.680 ^a	42.714 ^{ab}	17.907 ^a
		0.00	112.053 ^c	4.853 ^b	10.221 ^d	9.457 ^c
Zinnia rosso scarlatta	0	0.50	122.463 ^c	7.233 ^{ab}	34.497 ^{bc}	9.289 ^c
		0.75	149.283 ^{bc}	7.980 ^{ab}	34.309 ^{bc}	9.947 ^c
		0.00	177.400 ^{bc}	7.117 ^{ab}	16.415 ^d	9.668 ^c
	50	0.50	224.503 ^{abc}	12.087 ^{ab}	44.715 ^a	13.507 ^b
		0.75	242.060 ^{abc}	12.903 ^{ab}	43.239 ^{ab}	15.377 ^{ab}
	100	0.00	135.937 ^c	8.657 ^{ab}	17.800 ^d	9.830 ^c
		0.50	130.807 ^c	15.003 ^a	32.155 ^c	14.648 ^{ab}
		0.75	342.507 ^a	14.840 ^a	41.698 ^{ab}	17.112 ^{ab}

Discussion

Zinnia seedlings, when treated with varying concentrations of BA and ZnSO_4 , showed increases in several growth measurements presented in Tables 2, 3, and 4. Regarding the main impact of varying BA rates on plant growth parameters, show when the plants were sprayed with BA and give much higher vegetative growth traits than the control treatment. Spraying plants with BA at 100 mg l^{-1} gave the highest mean values of plant height, branches number per plant, leaf area, shoot fresh and dry weights per plant during the study for two cultivars compared to the control treatments. However, there was no significant difference recorded between the two concentrations of BA (50, and 100 mg l^{-1}) in branches number per plant and leaf area. In general, BA has the ability to promote cell division and elongation may account for its greater impact on vegetative development metrics. [25, 26], which leads to stimulation of primordial production and partially intermodal elongation on the apex [27], which is reflected in the increase of plant height. Cytokinins are essential for preventing or reducing apical dominance and promoting the release of axillary buds from apical dominance. [28, 29 ,30] which lead to an increase in branches number and leaves number per plant. The results obtained are comparable with those obtained by [25] on *Codiaeum variegatum* L. Cytokinin is a growth-control hormone that stimulates the production of proteins, cell division, expansion, and the mobilization of nutrients

[31]. Also, exogenous applications of BA are made to promote cell expansion which leads to increasing the leaf area [32]. The general increase in leaf area because of BA treatments agrees with the findings of [33] on *Helleborus Oriental*, [34] on *Echinacea Hybrids* and [35] on the aster plant

The positive effect of BA on shoot fresh and dry weights could be interpreted by the findings of [36] who believed that BA functions in promoting xylem differentiation and vascular strand development, which in turn led to more absorption of water and nutrients from the soil and. which was reflected in higher growth and may account for the greatest fresh weight also it could be attributed to the increment in plant height and branches number because of overcoming the apical dominance of the plant leading to the formation of more leaves. The effect on shoot fresh and dry weights was authenticated by may authors. [37] on *Polianthus tuberosa* and [38] on *Calandula officinalis* concluded that BA application increased the shoot fresh and dry weights. [39] reported that the maximum values of shoot fresh and dry weights of some ornamental plants resulted from BA at 25 mg⁻¹. These results obtained by [40] on Coroton plant, and [41] on gladiolus.

On the other hand, the effect of different rates of Zn spraying on plant growth parameters, show in Table (2, 3 and 4) indicated that spraying zinnia plants with different levels of Zn significantly increased vegetative growth parameters compared to control treatment. Spraying plants with Zn at 0.75 % gave the highest values of plant height, branch number per plant, leaf area, and shoot fresh and dry weights per plant during the study on the two cultivars compared to the control treatments. One of the reasons plants grow taller than untreated plants is because zinc has an advantageous impact on the biosynthesis of auxin hormones, which can promote cell division and elongation [42] reported that the positive effect of Zn on plant growth may be due to its roles in auxin biosynthesis through the amino acid tryptophan. Moreover, [43] and [44] demonstrated that the rise in internode length caused by the enlargement of protoplasm in cell wall material proportion and consequently, the resulting increase in cell size that showed up as intermodal elongation may be the main reason for an increase in shoot height after zinc treatment.

Also, increasing the chlorophyll content, leaf area, and another parameter by applying zinc sulfate can improve photosynthesis and enhance dry matter accumulation and lead to increased shoot dry weights, which will result in increasing the vegetative growth. Foliar zinc has significantly affected the leaf area which might be due the fact that zinc activated different types of enzymes which ultimately increased the leaf area of zinnia plant. These results are in line with [45]. The foliar application of zinc was significantly increasing development growth and size of spike of gladiolus plant [24]. Foliar zinc application significantly affected the leaves number also; zinc might be activated by different types of enzymes which increase the number of leaves and leaf area. These results are in line with [46] who reported that zinc application significantly increased plant height, number of leaves and leaf area of gladiolus plant.

CONCLUSION

In conclusion, the present study indicated that the growth parameters of two cultivars of zinnia plants were highly enhanced by spraying different concentrations of BA and varied levels of zinc. 100 mg⁻¹ is considered the best treatment to produce admirable vegetative parameters of zinnia plants under the environmental conditions of this study. As well as the zinc application levels increasing growth, based on the experimental results, the following conclusions were obtained: number of leaves, number of branches, number of internodes, leaf area (cm²) and plant height performed better with foliar Zn application of 0.50 and 0.75 percent combined with 100 mg⁻¹ **at BA**.

References

- [1]. Dole, J. M., & Wilkins, H. F. (1999). *Floriculture: Principles and species* Hall, Upper Saddle River, N.J., USA, 1048 p.
- [2]. Rouhi, V., Shiran, B., & Mohamadkhani, A. (2014). Effects of calcium chloride, gibberellin and benzyladenine on qualitative and quantitative characteristics and flower longevity of Zinnia (*Zinnia elegans* J.). *Journal of Horticultural Science*, 27(4), 444-452.
- [3]. Shiravand, D. (2011). *Design of green spaces with ornamental trees and shrubs*. Iran: Agriculture Training and Education Press.
- [4]. [4] Ziobro, K.S., Usova, K.A. (2016) The Use of Growth Regulators In The Cultivation Of Graceful Zinnia. New science: experience, tradition, innovation, v.591, n.2, p.4-8.
- [5]. Konstantinova, A.A. (2017) Characteristics of Flower Decoration In The City Of Onega. International Student Scientific Bulletin, v.5, p.92.
- [6]. Javid, Q. A., Abbasi, N. A., Hafiz, I. A., & Mughal, A. L. (2005). Performance of Zinnia (*Zinnia elegans*) "Dahlia Flowered" Crimson Shade by application of NPK fertilizer. *International Journal of Agriculture and Biology*, v.7, n.3, p.474-476.
- [7]. Loyola, C. E., Dole, J. M., & Dunning, R. (2019). North American specialty cut flower production and postharvest survey. *HortTechnology*, 29(3), 338-359.
- [8]. Kim, D. H., Doyle, M. R., Sung, S., & Amasino, R. M. (2009). Vernalization: winter and the timing of flowering in plants. *Annual Review of Cell and Developmental*, 25(1), 277-299.

[9]. [9] Safdar, A.W.; Tanveer, F.M.; Noor, U.N.M.; Niaz, A.W. (2016). Photoperiodic Effect on Vegetative Growth and Flower Quality of Zinnia (Zinnia elegans Jacq.). *Sarhad Journal of Agriculture*, 32(4). p.258-423.

[10]. Toscano, S., & Romano, D. (2021). Morphological, physiological, and biochemical responses of zinnia to drought stress. *Horticulturae*, 7(10), 362.

[11]. Marković, M., Šoštarić, J., Kojić, A., Popović, B., Bubalo, A., Bošnjak, D., & Stanisavljević, A. (2022). Zinnia (Zinnia elegans L.) and periwinkle (Catharanthus roseus (L.) G. Don) responses to salinity stress. *Water*, 14(7), 1066.

[12]. Yassin, M. Y., & Ismail, A. E. (1995). Effect of Zinnia elegans as a mix-crop along with tomato against Meloidogyne incognita and Rotylenchulus reniformis. *Der Tropenlandwirt-Journal of Agriculture in the Tropics and Subtropics*, 96(2), 221-225.

[13]. Abbasi, N. A., Zahoor, S., & Nazir, K. (2004). Effects of preharvest phosphorus and potassium fertilizers and postharvest AgNO₃ pulsing on the postharvest quality and shelf life of zinnia (Zinnia elegans cv. Blue point) cut flowers. *Int. J. Agri. Biol.*, 129-131.

[14]. Peng XiaoLi, P. X., Rao JingPing, R. J., & Zhang YanLong, Z. Y. (2007). Effect of exogenous salicylic acid on vase life of cut flowers of Prato lily and related physiological influence. *Acta Hort. Sinica*, 34(1): 189-192.

[15]. Yamaguchi, S., & Kamiya, Y. (2000). Gibberellin biosynthesis: its regulation by endogenous and environmental signals. *Plant and cell physiology*, 41(3), 251-257.

[16]. Chatsudthipong, V., & Muanprasat, C. (2009). Stevioside and related compounds: therapeutic benefits beyond sweetness. *Pharmacology & therapeutics*, 121(1), 41-54.

[17]. Riedell, W. E., Khoo, U., & Inglett, G. E. (1985). Effects of bioregulators on soybean leaf structure and chlorophyll retention. In: *Plant Growth Regulation*, Lake Alfred, Florida, Proceedings. Lake Alfred. 204-212

[18]. Dyer, D. J., Carlson, D. R., Cotterman, C. D., Sikorski, J. A., & Ditson, S. L. (1987). Soybean pod set enhancement with synthetic cytokinin analogs. *Plant Physiology*, 84(2), 240-243.

[19]. Hossain, M. S., Rahman, M. M., Rashid, M. R., Farid, A. T. M., Kaium, M. A., Ahmed, M., Alam, M. S. and Uddin, K. M. S. (2006). In "Handbook on Agro technology". 4th ed., Bangladesh Agricuklural Res. Inst., Gazipur1701, Bangladesh, 356-358

[20]. George, E. F., Hall, M. A., & De Klerk, G. J. (2008). Plant propagation by tissue culture 3rd Edition. *The Netherland, The Background Springer*, 65-175.

[21]. Cakmak, I. (2000). Tansley Review No. 111 Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *The New Phytologist*, 146(2), 185-205.

[22]. Khan, A., Ullah, F., Khan, M., Zeb, A., Khan, M., Ahmad, K., & Zainub, B. (2018). Response of Zinnia to foliar application of boron and zinc. *Science International*, 30, 133-139.

[23]. Mirshekari, B. (2012). Seed priming with iron and boron enhances germination and yield of dill (*Anethum graveolens*). *Turkish Journal of Agriculture and Forestry*, 36(1), 27-33.

[24]. Katiyar P., P. K., Chaturvedi, O. P., & Dheerendra Katiyar, D. K. (2012). Effect of foliar spray of zinc, calcium and boron on spike production of gladiolus cv. eurovision. 1(4): 334-338.

[25]. Mazher, A. A., Zaghloul, S. M., Mahmoud, S. A., & Siam, H. S. (2011). Stimulatory effect of kinetin, ascorbic acid and glutamic acid on growth and chemical constituents of *Codiaeum variegatum* L. plants. *American Eurasian J. Agric. & Environ. Sci.*, 10 (3): 318- 323.

[26]. Sadak, M. S., Dawood, M. G., Bakry, B. A., & El-Karamany, M. F. (2013). Synergistic effect of indole acetic acid and kinetin on performance, some biochemical constituents and yield of faba bean plant grown under newly reclaimed sandy soil. *World Journal of Agricultural Sciences*, 9(4), 335-344.

[27]. Kumari, S. (2017). Effect of Kinetin (6-FAP) and Cycocel (CCC) on growth, metabolism and cellular organelles in Pearl Millet (*Pennisetum glaucum*) under water stress. *Int. J. Curr. Microbiol. App. Sci*, 6(8), 2711-2719.

[28]. Tamas, I. A. (1995). Hormonal regulation of apical dominance. In *Plant hormones: physiology, biochemistry and molecular biology* (pp. 572-597). Dordrecht: Springer Netherlands. 572- 597.

[29]. Arteca, R. N. (1996). Plant growth substances: principles and applications. Springer Science & Business Media. Chapman & Hall, New York. 133: 67- 69.

[30]. Menaka, P., Rani, Y. A., Rao, K. N., Babu, P. H., & Ahamed, M. L (2018). Response of chickpea (*Cicer arietinum* L.) to foliar application of ethrel, kinetin and boron. *Int. J. Curr. Microbiol. App. Sci*, 7(11), 1653- 1660.

[31]. Bairwa, S., & Mishra, J. S. (2017). Effect of NAA, BA and kinetin on yield of African marigold (*Tagetes erecta* Linn.). *International Journal of Current Microbiology and Applied Sciences*, 6(6), 1236-1241.

[32]. Miller, C. O. (1956). Similarity of Some Kinetin and Red-Light Effects. *Plant Physiology*, 31(4), 318.

[33]. Henschke, M., Czuchaj, P. K., & Szczepaniak, S. J. (2015). The effect of benzyladenine and gibberellic acid on growth and flowering of Helleborus Orientalislam. *Bulgarian Journal of Agricultural Science*, 21(6), 1198-1203.

[34]. Mara, C. G. (2017). *Controlling growth in Echinacea Hybrids* (Doctoral dissertation, Ph. D. Thesis, Virginia Polytechnic Institute and state University).

[35]. Mohamed, Y. F. Y. (2017). Effect of some growth stimulants on growth, flowering and postharvest quality of aster (*Sympyotrichum novi-belgii* L.) cv. Purple Monarch. *Middle East J. Agric. Res.*, 6(2), 264-273.

[36]. Sorokin, H. P., & Thimann, K. V. (1964). The histological basis for inhibition of axillary buds in *Pisum sativum* and the effects of auxins and kinetin on xylem development. *Protoplasma*, 59(2), 326-350.

[37]. El-Sayed, A. A., Salem, M. A., & El-Maadawy, E. I. (1989). Effect of Gibberellic acid (GA3) and benzyladenine (BA) on *Polianthes tuberosa* L. *Journal of Agricultural Research, Tanta Univ. (Egypt)*, 15(2):301-311.

[38]. Menesi, F. A., Nofal, E. M. S., & El-Mahrouk, E. M. (1991). Effect of some growth regulators on *Calendula officinalis* L. *Egypt. J. Appl. Sci.*, 6, 1-15.

[39]. Menesi, F. A., Khalafalla, M. M., & Kandeel, Y. (1994). Effect of some growth regulators and calcium chloride on *Tagetes erecta*. The 1st Conf. Ornamental Hor. Fac. Agric. Kafr El-Sheikh, Tanta Univ., Egypt, 2:408-425.

[40]. Eid, R. A., & Abou-Leila, B. H. (2006). Response of croton plants to Gibberellic acid, benzyl adenine and ascorbic acid application. *World J. Agric. Sci* 2.2: 174-179.

[41]. Neetu, R. K., & Singh, A. K. (2016). Effect of different kinetin concentration on growth and flowering attributes in gladiolus cultivars. *Agricultural Science Digest-A Research Journal*, 36(4), 319-322.

[42]. Price, C. A. (1970). *Molecular approaches to plant physiology*. New York: McGraw-Hill. pp: 398.

[43]. El Bagoury, O. H., Mohamed, F. A., Sharabash, M. T. M., El Kholy, A. F., & Hashim, M. M. F. (1984). Response of soybean plant to zinc-chelate and soil moisture content [Egypt]. *Agric. Res. Rev.*, 26(413): 447-457.

[44]. Osman, A. S., Abido, Y. M. Y., & Allam, S. M. M. (2000). Response of soybean to phosphorus and zinc fertilization under irrigation regime. *Annals Agric. Sci.*, Ain Shams Univ., Cairo, 45(1): 229-238.

[45]. [Tariq, M., Hameed, S., Malik, K. A., & Hafeez, F. Y. (2007). Plant root associated bacteria for zinc mobilization in rice. *Pakistan Journal of Botany*, 39(1), 245.

[46]. Memon, S. A., Baloch, A. R., Muhammad Ayubbaloch, M. A., & Mahmooda Buriro, M. B. (2013). Effect of zinc sulphate and iron sulphate on the growth and flower production of gladiolus (*Gladiolus hortulanus*). *J. of Agric.* 9(6):1621-1630.

تأثير مستويات مختلفة من الزنك والبنزيل الأدينين عن طريق الرش على النمو الخضري لصنفين من نبات الزينيا *Zinnia elegans*.

ليلي شعبان محمد¹

نيروش سليمان حسن¹

بان جبرانيل خوشابا¹

اميره عبد الحميد يوسف²

¹قسم البستنة، كلية علوم الهندسة الزراعية، جامعة دهوك،إقليم كوردستان العراق.

²قسم الحدائق النباتية مركز البحوث الزراعية، جمهورية مصر العربية.

الخلاصة

أجريت هذه الدراسة في مشتل قسم البستنة، كلية علوم الهندسة الزراعية، جامعة دهوك. إقليم كردستان، العراق، لدراسة تأثير مستويات مختلفة من الزنك (0، 0.50، و 0.75) وتركيزات مختلفة من BA (0، 50 و 100) ملغم/لتر على بعض المعايير الخضرية لصنفين من نبات الزينيا. وكانت النتائج كما يلي: أظهرت القيم المتوسطة أن أكبر النباتات (90.556 سم) تم تسجيلها في صنف *Zinnia Lavonda*. كما أظهر نفس الصنف أقصى عدد من الفروع والوزن الطازج والجاف والكاروتين الكلي الذي بلغ (8,160 غ، 8,194 غ، 11.534 غ، 213.194 غ، 11.398 غ، 12.394 غ، 11.160 غ، 10.790 غ، 10.206 غ، 9.206 غ، 30.561 غ) على التوالي. بينما كانت تنتائج عدد الأوراق، وعدد السلاليات، ومساحة الورقة، والكلوروفيل الكلي، في صنف *Zinnia rosso scarlatta*، حيث بلغت أعلى النتائج (23.284)، (30.889)، (30.222)، (9.778)، (9.778)، (30.080)، (33.033)، (42.714)، (17.90)، (15.680)، (15.230) على التوالي. حول العوامل الثلاثة، أقصى ارتفاع للنباتات لصنفين *Zinnia rosso scarlatta* و *Zinnia Lavanda* ، وصلت القيم الأعلى إلى 101.411 سم عند رش النباتات بـ 50 ملغم/لتر-1 + 0.50% زنك على التوالي، مقارنة بمعاملة المقارنة التي بلغت 96.622 سم عند 0.50% زنك على التوالي، مقارنة بمعاملة المقارنة التي بلغت 77.856 سم على التوالي، بينما زاد عدد الأوراق وعدد الفروع وعدد العقد ومساحة الورقة (سم²) بشكل ملحوظ مقارنة معاملة المقارنة والمعاملات الأخرى، وصلت القيمة الرئيسية إلى (30.889) على التوالي، بينما زاد عدد الأوراق وعدد العقد ومساحة الورقة (سم²) على التوالي، عند رش صنف *Zinnia Lavonda* بـ 100 ملغم/لتر BA + 0.50% زنك. كما أن الوزن الطازج للنباتات، والوزن الجاف للنباتات، والكلوروفيل الكلي، والكاروتين الكلي أعطت أعلى القيم التي تم الوصول إليها (30.889)، (30.222)، (9.778)، (9.778)، (30.080)، (33.033)، (42.714)، (17.90)، (15.680)، (15.230) على التوالي، عند رش صنف *Zinnia* بـ 100 ملغم/لتر BA + 0.75% زنك.

الكلمات المفتاحية: مستويات الزنك، تركيزات (BA)، الرش، النمو الخضري، صنفين من نبات الزينيا.